

BRAIDS, THEIR PROPERTIES AND GENERALIZATIONS

V. V. VERSHININ

ABSTRACT. In the paper we give a survey on braid groups and subjects connected with them. We start with the initial definition, then we give several interpretations as well as several presentations of these groups. Burau presentation for the pure braid group and the Markov normal form are given next. Garside normal form and his solution of the conjugacy problem are presented as well as more recent results on the ordering and on the linearity of braid groups. Next topics are the generalizations of braids, their homological properties and connections with the other mathematical fields, like knot theory (via Alexander and Markov theorems) and homotopy groups of spheres.

CONTENTS

1. Introduction	2
2. Historical Remarks	2
3. Definitions and General Properties	4
3.1. Systems of n curves in three-dimensional space and braid groups	4
3.2. Braid groups and configuration spaces	6
3.3. Braid groups as automorphism groups of free groups and the word problem	6
3.4. Commutator subgroup and other presentations	7
3.5. Presentation of the pure braid group and Markov normal form	9
4. Garside Normal Form, Center and Conjugacy Problem	9
5. Ordering of Braids	12
6. Representations	13
6.1. Burau representation	13
6.2. Lawrence-Krammer representation	13
7. Generalizations of Braids	14
7.1. Configuration spaces of manifolds	14
7.2. Artin-Brieskorn braid groups	14
7.3. Braid groups of surfaces	16
7.4. Braid groups in handlebodies	16
7.5. Braids with singularities	18
8. Homological Properties	21
8.1. Configuration spaces and $K(\pi, 1)$ -spaces	21
8.2. Cohomology of pure braid groups	22
8.3. Homology of braid groups	23
9. Connections with the Other Domains	24

2000 *Mathematics Subject Classification.* Primary 20F36; Secondary 20F38, 57M.

Key words and phrases. Braid, presentation, pure braid group, representation, configuration space, Artin-Brieskorn group, word problem, Garside normal form, conjugacy problem, singular braid monoid, Markov Theorem.

The author was supported in part by the by CNRS-NSF grant No 17149, INTAS grant No 03-5-3251 and the ACI project ACI-NIM-2004-243 "Braids and Knots".

9.1. Markov Theorem	24
9.2. Homotopy groups of spheres and Makanin braids	25
References	26

1. INTRODUCTION

Braid groups describe intuitive concept of classes of continuous deformations of braids, which are collections of intertwining strands whose endpoints are fixed. Mathematically they can be considered from various points of view. The first intuitive approach is formalized naturally as isotopy classes of a collection of n connected curves (strings) in 3-dimensional space. This point of view is connected with the definition of braid group as the fundamental group of configuration space of n points on a plane. Also braids can be interpreted as a mapping class group of a punctured disc and as a subgroup of automorphism group of a free group (subsection 3.3).

The present survey is organized as follows. In Section 2 we make some historical remark. Definition and general properties are considered in Section 3. Configuration spaces appear in Subsection 3.2. Connections with groups of automorphisms of free groups are given in Subsection 3.3. Presentations of the braid group which appeared quite recently are observed in Subsection 3.4. Section 4 is devoted to F. A. Garside's classical work [91] and Section 5 to that of P. Dehornoy on ordering for braids. Representations and in particular linearity are discussed in Section 6. In Section 7 various generalization of braids are presented. Homological properties are observed in Section 8. In the last Section 9 we discuss connection with the knot theory given by the Alexander and Markov Theorems and with the homotopy groups of spheres.

The author is grateful to F. R. Cohen and Wu Jie for fruitful discussions on theory of braids and their applications. The author is also thankful to Vik. S. Kulikov who attracted his attention to the works of O. Zariski [201], [202] and to E. P. Volokitin for the help and advices on the presentation of the paper.

2. HISTORICAL REMARKS

Braids were rigorously defined by E. Artin [7] in 1925, although the roots of this natural concept are seen in the works of A. Hurwitz ([117], 1891), R. Fricke and F. Klein ([89], 1897) and even in the notebooks of K.-F. Gauss. E. Artin [7] gave the presentation of the braid group (see formulas (3.2) in Section 3) which is common now. Already in the book of Felix Klein [126] published in 1926 there appeared a chapter about braids. Essential topics about braids were also presented in the Reidemeister's Knotentheorie [176] published in 1932.

In 30-ies there appeared a series of papers of Werner Burau [48], [49], [50] where he in particular gave the presentation of the pure braid group (see subsection 3.5) and introduced the Burau representation (subsection 6.1). Wilhelm Magnus in his work [139] published in 1934 established relations between braid groups and the mapping class groups. At the same time there appeared the work of A. A. Markov [150], which together with Alexander Theorem [3] builds a bijection between links and equivalence classes of braids. It became an essential ingredient in study of links and knots (in the work of V. F. R. Jones [121], for example). In 1936-37 were published the works of O. Zariski [201], [202], where he discovered connections between braid groups and the fundamental group of the complement of discriminant of the general polynomial

$$f_n(t) = a_0 t^n + a_1 t^{n-1} + \dots + a_{n-1} t + a_n,$$

a point of view later rediscovered by V. I. Arnold [5]. Zariski also understood connections between braids and configuration spaces, gave the presentation of the braid group of the sphere, and studied the braid groups of Riemann surfaces. Amazingly and unfortunately these works of Zariski were not noticed by the specialists on braids and are not mentioned even in books and papers where the presentations of braids of surfaces are discussed.

In the beginning of 60-ies R. Fox and L. Neuwirth [86] and E. Fadell and L. Neuwirth [80] studied configuration spaces which turned out to be $K(\pi, 1)$ -spaces and so give natural geometrical model of the classifying spaces for the braid groups. Later, V. I. Arnold [5] in this direction proved the first results on cohomologies of braids. The motivation for his study was a connection (which he discovered) with the problem of representing algebraic functions in several variables by superposing algebraic functions in fewer variables. Also, in 1969 V. I. Arnold completely described the cohomologies of pure braid groups [4].

In 1969 there appeared the publication of F. A. Garside's work [91] where he suggests a new normal form of elements in the braid group and with its help gives a new solution of the word problem and also solves the conjugacy problem. In 1968 was published a two-page note of G. S. Makanin [143] where he sketches his algorithm for the solution of the conjugacy problem. The complete publication of Makanin's work didn't appear (as far as the author is aware).

In 70-ies the study of cohomologies of braids was continued independently and by different methods by D. B. Fuks [90] who determined mod 2 cohomologies, and F. R. Cohen [54], [55], [56] who described these homologies with coefficients in $\mathbb{Z}$ and in $\mathbb{Z}/p$ as modules over the Steenrod algebra.

In 1984-85 independently N. V. Ivanov [118] and J. McCarthy [152] proved the "Tits-alternative" for the mapping class groups of surfaces and as a consequence it is true for the braid groups. Namely, they proved that every subgroup of the mapping class group either contains an abelian subgroup of finite index, or contains a non-abelian free group.

The question of whether braid groups are linear attracted significant attention. It was realized that the Burau representation is faithful for Br_3 [92], [142]. Then after a long break in 1991 J. A. Moody [154] proved that Burau representation is unfaithful for $n \geq 9$. This bound was improved to $n \geq 6$ by D. D. Long and M. Paton [138] and to $n = 5$ by S. Bigelow [28]. In 1999-2000 there appeared preprints of papers of D. Krammer [128], [129] and S. Bigelow [29] who proved that Br_n is linear for all n (using the other representation).

At the begining of nineties P. Dehornoy [67], [68], [69] proved that there exists a left order in braid groups.

Interesting generalizations of braids were introduced in the work of E. Brieskorn [42]. The configuration space can also be considered as the orbit space of the complement of the complexification of the arrangement of hyperplanes corresponding to the Coxeter group $A_{n-1} = \Sigma_n$. Generalizing this approach for any finite Coxeter group, E. Brieskorn defined the so-called generalized braid groups which are also called Artin groups.

Another way of generalization is to consider braid groups in 3-manifolds, possibly with a boundary. The simplest examples are braid groups in handlebodies. A. B. Sossinsky [183] was the first who studied them. Such a group can be interpreted as the fundamental group of the configuration space of a plane without g points where g is the genus of the handlebody. The generalized braid group of type C is isomorphic to the braid group in the solid torus.

Under the influence of the theory of Vassiliev-Goussarov (finite - type) invariants singular braids were introduced. The corresponding algebraic structures are the Baez-Birman monoid




FIGURE 3.1.




FIGURE 3.2.

[10], [33] and the braid-permutation group by R. Fenn, R. Rimányi and C. Rourke [83], [84]. Various properties of these objects were studied [85], [203], [93], [94], [66], [119], [62], [104].

3. DEFINITIONS AND GENERAL PROPERTIES

3.1. Systems of n curves in three-dimensional space and braid groups. First of all as it was already mentioned braids naturally arise as objects in 3-space. Let us consider two parallel planes P_0 and P_1 in $\mathbb{R}^3$, which contain two ordered sets of points $A_1, \dots, A_n \in P_0$ and $B_1, \dots, B_n \in P_1$. These points are lying on parallel lines L_A and L_B respectively. The space between the planes P_0 and P_1 we denote by Π . Suppose that the point B_i is lying under the point A_i , as a result of the orthogonal projection of the plane P_0 onto the plane P_1 . Let us connect the set of points $A_1, \dots, A_n$ with the set of points $B_1, \dots, B_n$ by simple non-intersecting curves $C_1, \dots, C_n$ lying in the space Π and such that each curve meets only once each parallel plane P_t lying in the space Π (see Figure 3.1). This object is called a *braid* and the curves are called the *strings* of a braid. Usually braids are depicted by projections on the plane passing through the lines L_A and L_B . This projection is supposed to be in general position so that there is only finite number of double points of intersection which are lying on pairwise different levels and intersections are transversal. The simplest braid σ_i (Fig. 3.2) corresponds to the transposition $(i, i+1)$.

Let us introduce the following equivalence relation on the set of all braids with n strings and with fixed P_0, P_1, A_i and B_i . It is defined by homeomorphisms $h : \Pi \rightarrow \Pi$, identical on $P_0 \cup P_1$ and such that $h(P_t) = P_t$. Braids β and β' are equivalent if there exists such a homeomorphism h that $h(\beta) = \beta'$. On the set Br_n of equivalence classes under the considered relation the structure of a group introduces as follows. We put a copy Π' of the domain Π under the Π in such a way that P'_0 coincides with P_1 and each A_i coincides with B_i and we glue braids β and




FIGURE 3.3.

β' . This gluing gives a composition of braids $\beta\beta'$ (Fig. 3.3). Unit element is the equivalence class containing a braid of n parallel intervals, the braid β^{-1} inverse to β is defined by reflection of β with respect to the plane $P_{1/2}$. A string C_i of a braid β connects the point A_i with the point B_{k_i} defining the permutation S^β . If this permutation is identical then the braid β is called *pure*. The map $\beta \rightarrow S^\beta$ defines an epimorphism τ_n of the braid group Br_n on the permutation group Σ_n with the kernel consisting of all pure braids:

$$(3.1) \quad 1 \rightarrow P_n \rightarrow Br_n \xrightarrow{\tau_n} \Sigma_n \rightarrow 1.$$

The following presentation of the braid group Br_n with generators σ_i , $i = 1, \dots, n-1$ and two types of relations:

$$(3.2) \quad \begin{cases} \sigma_i \sigma_j &= \sigma_j \sigma_i, \text{ if } |i-j| > 1, \\ \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1} \end{cases}$$

is the algebraic expression of the fact that any isotopy of braids can be broken down into “elementary moves” of two types that correspond to two types of relations.

If we add a vertical interval to the system of curves on Figure 3.1 we can get a canonical inclusion j_n of the group Br_n into the group Br_{n+1}

$$j_n : Br_n \rightarrow Br_{n+1}.$$

If the symmetric group Σ_n is given by its canonical presentation with generators s_i , $i = 1, \dots, n-1$ and relations:

$$(3.3) \quad \begin{cases} s_i s_j &= s_j s_i, \text{ if } |i-j| > 1, \\ s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1} \\ s_i^2 &= 1, \end{cases}$$

then the homomorphism τ_n is given by the formula

$$\tau_n(\sigma_i) = s_i, \quad i = 1, \dots, n-1.$$

It is possible to consider braids as classes of equivalence of *braid diagrams* which are generic projections of three dimensional braids on a plane. The classes of equivalence are defined by the *Reidemeister moves* depicted at Figure 3.4.




FIGURE 3.4.



FIGURE 3.5.

3.2. Braid groups and configuration spaces. If we look at the Figure 3.1, then this picture can be interpreted as a graph of a loop in the *configuration space* of n points on a plane, that is the space of unordered sets of n points on a plane, see Figure 3.5. So, it is possible to interpret the braid group as the fundamental group of the configuration space. Formally it is done as follows. The symmetric group Σ_m acts on the Cartesian power $(\mathbb{R}^2)^m$ of the space $\mathbb{R}^2$:

$$(3.4) \quad w(y_1, \dots, y_m) = (y_{w^{-1}(1)}, \dots, y_{w^{-1}(m)}), \quad w \in \Sigma_m.$$

Denote by $F(\mathbb{R}^2, m)$ the space of m -tuples of pairwise different points in $\mathbb{R}^2$:

$$F(\mathbb{R}^2, m) = \{(p_1, \dots, p_m) \in (\mathbb{R}^2)^m : p_i \neq p_j \text{ for } i \neq j\}.$$

This is the space of regular points of our action. We call the orbit space of this action $B(\mathbb{R}^2, m) = F(\mathbb{R}^2, m)/\Sigma_m$ the *configuration space of n points on a plane*. The braid group Br_m is the fundamental group of configuration space

$$Br_m = \pi_1(B(\mathbb{R}^2, m)).$$

The pure braid group P_m is the is the fundamental group of the space $F(\mathbb{R}^2, m)$. The covering

$$p : F(\mathbb{R}^2, m) \rightarrow B(\mathbb{R}^2, m)$$

defines the exact sequence:

$$(3.5) \quad 1 \rightarrow \pi_1(F(\mathbb{R}^2, m)) \xrightarrow{p_*} \pi_1(B(\mathbb{R}^2, m)) \rightarrow \Sigma_n \rightarrow 1,$$

which is equivalent to sequence (3.1).

It can be used for proving the canonical presentation of the braid group (3.2) as it is done, for example in the book of J. Birman [32].

Such considerations were made by R. Fox and L. Neuwirth [86].

3.3. Braid groups as automorphism groups of free groups and the word problem. Another important approach to the braid group bases on the fact that this group may be considered as a subgroup of the automorphism group of a free group.

Let F_n be the free group of rank n with the set of generators $\{x_1, \dots, x_n\}$. Assume further that $\text{Aut } F_n$ is the automorphism group of F_n .




FIGURE 3.6.

We have the standard inclusions of the symmetric group Σ_n and the braid group Br_n into $\text{Aut } F_n$. For the braid group it may be described as follows. Let $\bar{\sigma}_i \in \text{Aut } F_n, i = 1, 2, \dots, n-1$, be given by the formula which describes its action on generators:

$$(3.6) \quad \begin{cases} x_i & \mapsto x_{i+1}, \\ x_{i+1} & \mapsto x_{i+1}^{-1} x_i x_{i+1}, \\ x_j & \mapsto x_j, j \neq i, i+1. \end{cases}$$

Let us define a map ν of the generators $\sigma_i, i = 1, \dots, n-1$ of the braid group Br_n to these automorphisms:

$$(3.7) \quad \nu(\sigma_i) = \bar{\sigma}_i.$$

Theorem 3.1. *Formulas (3.7) define correctly a homomorphism*

$$\nu : Br_n \rightarrow \text{Aut } F_n.$$

which is a monomorphism.

Theorem 3.1 gives a solution of the word problem for the braid groups. It was done first by E. Artin [7].

The free group F_n is a fundamental group of a disc D_n without n points and the generator x_i corresponds to a loop going around the i -th point. The braid group Br_n is the mapping class group of a disc D_n with its boundary fixed [32] and so it acts on the fundamental group of D_n . This action is described by the formulas (3.6) where x_i correspond to the canonical loops on D_n which form the generators of the fundamental group. Geometrically this action is depicted in the Figure 3.6.

3.4. Commutator subgroup and other presentations. Let us define a homomorphism from braid group to integers by taking the sum of exponents of the entries of the generators σ_i in the expression of any element of the group through these canonical generators:

$$\deg : Br_n \rightarrow \mathbb{Z}, \deg(b) = \sum_j m_j, \text{ where } b = (\sigma_{i_1})^{m_1} \dots (\sigma_{i_k})^{m_k}.$$

Proposition 3.1. *The homomorphism*

$$\deg : Br_n \rightarrow \mathbb{Z}$$

gives the abelianization of the braid group and the commutator subgroup Br'_n is characterized by the condition

$$b \in Br'_n \text{ if and only if } \deg(b) = 0.$$




FIGURE 3.7.

Proof. Let $a : Br_n \rightarrow A$ be a homomorphism to any other abelian group A , then from the relations (3.2) we have:

$$a(\sigma_i)a(\sigma_{i+1})a(\sigma_i) = a(\sigma_{i+1})a(\sigma_i)a(\sigma_{i+1}).$$

the commutativity of A gives that $a(\sigma_{i+1}) = a(\sigma_i)$. This means that the homomorphism $\deg$ is universal. $\square$

Of course, there exist another presentations of the braid group. Let

$$\sigma = \sigma_1\sigma_2 \dots \sigma_{n-1},$$

then the group Br_n is generated by σ_1 and σ because

$$\sigma_{i+1} = \sigma^i \sigma_1 \sigma^{-i}, \quad i = 1, \dots, n-2.$$

The relations for the generators σ_1 and σ are the following

$$(3.8) \quad \begin{cases} \sigma_1 \sigma^i \sigma_1 \sigma^{-i} &= \sigma^i \sigma_1 \sigma^{-i} \sigma_1 \quad \text{for } 2 \leq i \leq n/2, \\ \sigma^n &= (\sigma \sigma_1)^{n-1}. \end{cases}$$

This was observed by Artin in the initial paper [7].

An interesting series of presentations was given by V. Sergiescu [182]. For every planar graph he constructed a presentation of the group Br_n , where n is the number of vertices of the graph, with generators corresponding to edges and relations reflecting the geometry of the graph. Artin's presentation in this context corresponds to the graph consisting of the interval from 1 to n with the natural numbers (from 1 to n) as vertices and with segments between them as edges. For generalizations of braids graph presentations of these type were considered by P. Bellingeri and V. Vershinin [17, 21]

J. S. Birman; K. H. Ko; S. J. Lee [35] introduced the presentation with the generators a_{ts} with $1 \leq s < t \leq n$ and relations

$$(3.9) \quad \begin{cases} a_{ts}a_{rq} &= a_{rq}a_{ts} \quad \text{for } (t-r)(t-q)(s-r)(s-q) > 0, \\ a_{ts}a_{sr} &= a_{tr}a_{ts} = a_{sr}a_{tr} \quad \text{for } 1 \leq r < s < t \leq n. \end{cases}$$

The generators a_{ts} are expressed by the canonical generators σ_i in the following form:

$$(3.10) \quad a_{ts} = (\sigma_{t-1}\sigma_{t-2} \dots \sigma_{s+1})\sigma_s(\sigma_{s+1}^{-1} \dots \sigma_{t-2}^{-1}\sigma_{t-1}^{-1}) \quad \text{for } 1 \leq s < t \leq n.$$

Geometrically the generators $a_{s,t}$ are depicted in Figure 3.7.

The set of generators for braid groups was even enlarged in the work of Jean Michel [153] as follows. Let $|| : \Sigma_n \rightarrow \mathbb{Z}$ be the length function on the symmetric group with respect to

the generators s_i : for $x \in \Sigma_n$, $|x|$ is the smallest natural number k such that x is a product of k elements of the set $\{s_1, \dots, s_{n-1}\}$. It is known ([41], Sect. 1, Ex. 13(b)) that two minimal expressions for an element of Σ_n are equivalent by using only of the relations (3.2). This implies that the canonical projection $\tau_n : Br_n \rightarrow \Sigma_n$ has a unique set-theoretic section $r : \Sigma_n \rightarrow Br_n$ such that $r(s_i) = \sigma_i$ for $i = 1, \dots, n-1$ and $r(xy) = r(x)r(y)$ whenever $|xy| = |x| + |y|$. Then the group Br_n admits a presentation by generators $\{r(x) \mid x \in \Sigma_n\}$ and relations $r(xy) = r(x)r(y)$ for all $x, y \in \Sigma_n$ such that $|xy| = |x| + |y|$.

3.5. Presentation of the pure braid group and Markov normal form. Let $f(y_1, \dots, y_m)$ be a word with (possibly empty) entries of y_i^ϵ , where y_i are some letters and ϵ may be ± 1 . If y_i are elements of a group G then $f(y_1, \dots, y_m)$ will be considered as the corresponding element of G .

Let us define the elements $s_{i,j}$, $1 \leq i < j \leq m$, of the braid group Br_m by the formula:

$$s_{i,j} = \sigma_{j-1} \dots \sigma_{i+1} \sigma_i^2 \sigma_{i+1}^{-1} \dots \sigma_{j-1}^{-1}.$$

These elements satisfy the following Burau relations ([48], [151], see also [135]):

$$(3.11) \quad \begin{cases} s_{i,j} s_{k,l} &= s_{k,l} s_{i,j} \text{ for } i < j < k < l \text{ and } i < k < l < j, \\ s_{i,j} s_{i,k} s_{j,k} &= s_{i,k} s_{j,k} s_{i,j} \text{ for } i < j < k, \\ s_{i,k} s_{j,k} s_{i,j} &= s_{j,k} s_{i,j} s_{i,k} \text{ for } i < j < k, \\ s_{i,k} s_{j,k} s_{j,l} s_{j,k}^{-1} &= s_{j,k} s_{j,l} s_{j,k}^{-1} s_{i,k} \text{ for } i < j < k < l. \end{cases}$$

W. Burau and later A. A. Markov proved that the elements $s_{i,j}$ with the relations (3.11) give a presentation of the pure braid group P_m [151]. The following formula is a consequence of the Burau relations and also belongs to A. A. Markov:

$$(3.12) \quad [s_{i,l}, s_{j,k}^\epsilon] = f(s_{1,l}, \dots, s_{l-1,l}), \quad \epsilon = \pm 1, \quad k < l.$$

Let us define the elements $\sigma_{k,l}$, $1 \leq k \leq l \leq m$ by the formulas

$$\begin{aligned} \sigma_{k,k} &= e, \\ \sigma_{k,l} &= \sigma_k^{-1} \dots \sigma_{l-1}^{-1}. \end{aligned}$$

Let P_m^k be the subgroup of P_m generated by the elements $s_{i,j}$ with $k < j$.

Theorem 3.2. (A. A. Markov) *(i) Every element of the group Br_m can be uniquely written in the form*

$$(3.13) \quad f_m(s_{1,m}, \dots, s_{m-1,m}) \dots f_j(s_{1,j}, \dots, s_{j-1,j}) \dots f_2(s_{1,2}) \sigma_{i_m,m} \dots \sigma_{i_j,j} \dots \sigma_{i_2,2}.$$

(ii) The factor group P_m^k / P_m^{k-1} is the free group on free generators $s_{i,k+1}$, $1 \leq i \leq k$.

The form (3.13) is called the *Markov normal form*, it also gives the solution of the word problem for the braid groups.

4. GARSIDE NORMAL FORM, CENTER AND CONJUGACY PROBLEM

Essential role in Garside work [91] plays the monoid of *positive* braids Br_n^+ , that is the monoid which has a presentation with generators σ_i , $i = 1, \dots, n$ and relations (3.2). In other words each element of this monoid can be represented as a word on the elements σ_i , $i = 1, \dots, n$ with no entrances of σ_i^{-1} . Two positive words A and B in the alphabet $\{\sigma_i, (i = 1, \dots, n-1)\}$ will be said to be *positively equal* if they are equal as elements of Br_n^+ . In this case we shall write $A \doteq B$.




FIGURE 4.1.

First of all Garside proves the following statement.

Proposition 4.1. *In Br_n^+ for $i, k = 1, \dots, n-1$, given $\sigma_i A \doteq \sigma_k B$, it follows that*

if $k = i$, then $A \doteq B$,

if $|k - i| = 1$, then $A \doteq \sigma_k \sigma_i Z$, $B \doteq \sigma_i \sigma_k Z$ for some Z ,

if $|k - i| \geq 2$, then $A \doteq \sigma_k Z$, $B \doteq \sigma_i Z$ for some Z .

The same is true for the right multiples of σ_i .

Corollary 4.1. *If $A \doteq P$, $B \doteq Q$, $AXB \doteq PYQ$, ($L(A) \geq 0$, $L(B) \geq 0$), then $X \doteq Y$. That is, monoid Br_n^+ is left and right cancellative.*

Garside's *fundamental word* Δ in the braid group Br_{n+1} is defined by the formula:

$$\Delta = \sigma_1 \dots \sigma_n \sigma_1 \dots \sigma_{n-1} \dots \sigma_1 \sigma_2 \sigma_1.$$

If we use Garside's notation $\Pi_t \equiv \sigma_1 \dots \sigma_t$, then $\Delta \equiv \Pi_{n-1} \dots \Pi_1$.

For a positive word W in σ_i , $i = 1, \dots, n$ we say that Δ is a *factor* of W or simply W contains Δ , if $W \doteq A\Delta B$ with A and B being arbitrary positive words, probably empty. If W does not contain Δ we shall say W is *prime to* Δ .

Garside's transformation of words $\mathcal{R}$ is defined by the formula

$$\mathcal{R}(\sigma_i) \equiv \sigma_{n-i}.$$

This gives the automorphism of Br_n and the positive braid monoid Br_n^+ .

Proposition 4.2. *In Br_n*

$$\sigma_i \Delta \doteq \Delta \mathcal{R}(\sigma_i).$$

Geometrically this commutation is shown on Figure 4.1 ($\Delta \sigma_3 = \sigma_1 \Delta$).

Proposition 4.3. *If W is an arbitrary positive word in Br_n^+ such that either*

$$W \doteq \sigma_1 A_1 \doteq \sigma_2 A_2 \doteq \dots \doteq \sigma_{n-1} A_{n-1},$$

or

$$W \doteq B_1 \sigma_1 \doteq B_2 \sigma_2 \doteq \dots \doteq B_{n-1} \sigma_{n-1},$$

then $W \doteq \Delta Z$ for some Z .

Proposition 4.4. *The canonical homomorphism*

$$Br_n^+ \rightarrow Br_n$$

is a monomorphism.

Among positive words on the alphabet $\{\sigma_1 \dots \sigma_n\}$ let us introduce a lexicographical ordering with the condition that $\sigma_1 < \sigma_2 < \dots < \sigma_n$. For a positive word W the *base* of W is the smallest positive word which is positively equal to W . The base is uniquely determined. If a positive word A is prime to Δ , then for the base of A the notation $\overline{A}$ will be used.

Theorem 4.1. (F. A. Garside) *Every word W in Br_{n+1} can be uniquely written in the form $\Delta^m \overline{A}$, where m is an integer.*

The form of a word W established in this theorem we call the *Garside left normal form* and the index m we call the *power* of W . The same way the *Garside right normal form* is defined and the corresponding variant of Theorem 4.1 is true. The Garside normal form also gives a solution to the word problem in the braid group.

Theorem 4.2. (F. A. Garside) *The necessary and sufficient condition that two words in Br_{n+1} are equal is that their Garside normal forms are identical.*

Garside normal form for the braid groups was precised in the subsequent works of S. I. Adyan [1], W. Thurston [78], E. El-Rifai and H. R. Morton [76]. Namely, there was introduced the *left-greedy form* (in the terminology of W. Thurston [78])

$$\Delta^t A_1 \dots A_k,$$

where A_i are the successive possible longest *fragments of the word Δ* (in the terminology of S. I. Adyan [1]) or *positive permutation braids* (in the terminology of E. El-Rifai and H. R. Morton [76]). Certainly, the same way the *right-greedy form* is defined. With the help of this form it proved that the braid group is biautomatic.

The center of the braid group was firstly found by W.-L. Chow [53]. Namely, as it follows from the presentation of braid groups with two generators σ_1 and σ and relations (3.8) given in the subsection 3.1 the element σ^n commutes with $\sigma\sigma_1$ and so with σ_1 . Chow proved that it generates the center. Garside normal form gives an elegant proof of the following theorem.

Theorem 4.3. (i) *When $n = 1$ the center of the group Br_{n+1} is generated by Δ .*

(ii) *When $n > 1$ the center of the group Br_{n+1} is generated by Δ^2 .*

Let α be a positive word such that $\Delta \doteq \alpha X$, where X is an arbitrary positive word, probably empty. For any word W in Br_{n+1} , the word $\alpha^{-1}W\alpha$, reduced to Garside normal form is called an α -*transformation* of W .

For any word W in Br_{n+1} with the Garside normal form $\Delta^m \overline{A} \equiv W_1$ consider the following chains of α -transformations: take all the α -transformations of W_1 and let those which are of power $\geq m$ and which are distinct from each other be $W_2, W_3, \dots, W_t$. Now repeat the process

for each of the words $W_2, W_3, \dots, W_t$ in turn, denoting successively by $W_{t+1}, W_{t+2}, \dots$, any new words occurring, the condition being always that each new word must be of power $\geq m$. Continue to repeat the process for every new distinct word arising, as the sequence $W_1, W_2, W_{t+2}, \dots$, expands.

Proposition 4.5. *The set $W_1, W_2, W_{t+2}, \dots$, is finite.*

Suppose that in the set $W_1, W_2, W_{t+2}, \dots$, the highest power reached is s and that the words of power s form the subset $V_1, V_2, \dots$. Then this set $V_1, V_2, \dots$ is called the *summit set* of W .

Theorem 4.4. (F. A. Garside) *Two elements A and B of the group Br_{n+1} are conjugate if and only if their summit sets are identical.*

J. S. Birman, K. H. Ko and S. J. Lee considered the word $\delta = a_{n(n-1)} \cdots a_{32}a_{21} = \sigma_{n-1} \cdots \sigma_2\sigma_1$, as a fundamental in their system of generators and proved that every element in Br_n has a representative $W = \delta^j A_1 A_2 \cdots A_k$ with positive A_i in a unique way in some sense. Based on this form they gave an algorithm for the word problem in B_n which runs in time (n^2) for a given word of length m .

5. ORDERING OF BRAIDS

A group G is said *totally* (or *linearly*) *left* (correspondingly *right*) *ordered* if it has a total order $<$ invariant by left (right) multiplication, i.e. if $a < b$, then $ca < cb$ for any $c \in G$. If this order is also invariant by right (left) multiplication, then the group G is called *ordered*.

For any left ordered group G denote by P the set of positive elements $\{x \in G : x > 1\}$, then the set of negative elements is defined by the formula: $P^{-1} = \{x \in G : x \in P\}$. The total character of an order on G is expressed by the partition

$$G = P \coprod \{1\} \coprod P^{-1}.$$

The invariance of multiplication is expressed by the inclusion $P^2 \subset P$, where P^2 is formed by products of couples of elements of P . Conversely, if there exist a subset P of a group G with the properties:

$$G = P \coprod \{1\} \coprod P^{-1}, \quad P^2 \subset P,$$

then G is left ordered by the order defined by: $x < y$ if and only if $x^{-1}y \in P$. A group G then is ordered if and only if $xPx^{-1} \subset P$ for all $x \in G$.

Let $i \in \{1, \dots, n\}$ and a word w on the alphabet $\{\sigma_1, \dots, \sigma_n\}$ is expressed in the form

$$w_0 \sigma_i w_1 \sigma_i \dots \sigma_i w_r,$$

where the subwords $w_0, \dots, w_r$ are the words on the letters $\sigma_j^{\pm 1}$ with $j > i$. Then such a word is called σ_i -*positive*. This means that all entries of $\sigma_i^{\pm 1}$ in the word w with i minimal must be positive. If all such entries are negative then a word w is called σ_i -*negative*. A braid of Br_{n+1} is called σ_i -*positive* (σ_i -*negative*) if there exists its expression as a word on the standard generators which is σ_i -*positive* (σ_i -*negative*). A braid is called σ -*positive* (σ -*negative*) if it exists a number i , such that it is σ_i -*positive* (σ_i -*negative*).

Theorem 5.1. (P. Dehornoy) *Every braid in Br_{n+1} different from 1 is either σ -positive or σ -negative.*

Corollary 5.1. *For all n the braid group Br_{n+1} is left ordered.*

6. REPRESENTATIONS

6.1. Burau representation. Let us map the generators of the braid group Br_n to the following elements of the group $GL_n\mathbb{Z}[t, t^{-1}]$

$$(6.1) \quad \sigma_i \mapsto \begin{pmatrix} E_{i-1} & 0 & 0 & 0 \\ 0 & 1-t & t & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & E_{n-i-1} \end{pmatrix},$$

where E_i is the unit $i \times i$ matrix. The formula (6.1) defines correctly the representation of the braid group in $GL_n\mathbb{Z}[t, t^{-1}]$:

$$r : Br_n \rightarrow GL_n\mathbb{Z}[t, t^{-1}],$$

which is called *Burau representation* [50].

Theorem 6.1. *Burau representation is faithful for $n = 3$.*

Theorem 6.2. (J. A. Moody; D. D. Long and M. Paton; S. Bigelow) *Burau representation is unfaithful for $n \geq 5$.*

The case $n = 4$ remains open.

6.2. Lawrence-Krammer representation. Consider the ring $K = \mathbb{Z}[q^{\pm 1}, t^{\pm 1}]$ of Laurent polynomials in two variables q, t , and the free K -module

$$V = \bigoplus_{1 \leq i < j \leq n} K x_{i,j}.$$

For $k \in \{1, 2, \dots, n-1\}$, define the action of the braid generators σ_k on the basis of V by the formula:

$$(6.2) \quad \sigma_k(x_{i,j}) = \begin{cases} x_{i,j}, & k < i-1 \text{ or } j < k; \\ x_{i-1,j} + (1-q)x_{i,j}, & k = i-1; \\ tq(q-1)x_{i,i+1} + qx_{i+1,j}, & k = i < j-1; \\ tq^2x_{i,j}; & k = i = j-1; \\ x_{i,j} + tq^{k-i}(q-1)^2x_{k,k+1}, & i < k < j-1; \\ x_{i,j-1} + tq^{j-i}(q-1)x_{j-1,j}, & k = j-1; \\ (1-q)x_{i,j} + qx_{i,j+1}, & k = j. \end{cases}$$

Direct computation shows that this defines a representation

$$\rho_n : Br_n \rightarrow GL(V),$$

which was firstly defined by R. Lawrence [134] in topological terms and in the explicit form (6.2) by D. Krammer [129].

Theorem 6.3. (S. Bigelow [29], D. Krammer [129]) *The representation*

$$\rho_n : Br_n \rightarrow GL(V)$$

is faithful for all $n \geq 1$.

Remark 6.1. *Actually, S Bigelow [29] proved this theorem for the representation ρ_n characterized in homological terms and D. Krammer [129] proved the following. Let $K = \mathbb{R}[t^{\pm 1}]$, $q \in \mathbb{R}$, and $0 < q < 1$. Then the representation ρ_n defined by (6.2) is faithful for all $n \geq 1$. This result implies Theorem 6.3: if a representation over $\mathbb{Z}[q^{\pm 1}, t^{\pm 1}]$ becomes faithful after assigning a real value to q , then it is faithful itself.*

M. G. Zinno [204] established connection between Birman-Murakami-Wenzl algebra [40], [156] and the Lawrence-Krammer representation. Namely, he proved that the Lawrence-Krammer representation is identical to the irreducible representations of the Birman-Murakami-Wenzl algebra parametrized by Young diagrams of shapes $(n-2)$ and (1^{n-2}) . This means that the Young diagram in the case considered consists of one row (respectively of one column) only, with $n-2$ boxes. It follows that Lawrence-Krammer representation is irreducible.

7. GENERALIZATIONS OF BRAIDS

7.1. Configuration spaces of manifolds. The notion of configuration space of Subsection 3.2 can be naturally generalized for a configuration space of a manifold as follows. Let Y be a connected topological manifold and let W be a finite group acting on Y . A point $y \in Y$ is called *regular* if its stabilizer $\{w \in W : wy = y\}$ is trivial, i.e., consists only of the unit of the group W . The set $\tilde{Y}$ of all regular points is open. Suppose that it is connected and nonempty. The subspace $ORB(Y, W)$ of the space of all orbits $Orb(Y, W)$ consisting of the orbits of all regular points is called the *space of regular orbits*. There is a free action of W on $\tilde{Y}$ and the projection $p : \tilde{Y} \rightarrow \tilde{Y}/W = ORB(Y, W)$ defines a covering. Let us consider the initial segment of the long exact sequence of this covering:

$$(7.1) \quad 1 \rightarrow \pi_1(\tilde{Y}, y_0) \xrightarrow{p_*} \pi_1(ORB(Y, W), p(y_0)) \rightarrow W \rightarrow 1.$$

The fundamental group $\pi_1(ORB(Y, W), p(y_0))$ of the space of regular orbits is called the *braid group of the action of W on Y* and is denoted by $Br(Y, W)$. The fundamental group $\pi_1(\tilde{Y}, y_0)$ is called the *pure braid group of the action of W on Y* and is denoted by $P(Y, W)$. The spaces $\tilde{Y}$ and $ORB(Y, W)$ are path connected, so the pair of these groups is defined uniquely up to isomorphism and we may omit mentioning the base point y_0 in the notations.

For any space Y the symmetric group Σ_m acts on the Cartesian power Y^m of the space Y by the formulas (3.4). We denote by $F(Y, m)$ the space of m -tuples of pairwise different points in Y :

$$F(Y, m) = \{(p_1, \dots, p_m) \in Y^m : p_i \neq p_j \text{ for } i \neq j\}.$$

This is the space of regular points of this action. In the case when Y is a connected topological manifold M without boundary and $\dim M \geq 2$, the space of regular orbits $ORB(M^m, \Sigma_m)$ is open, connected and nonempty. We call $ORB(M^m, \Sigma_m)$ the *configuration space of the manifold M* and denote by $B(M, m)$. The braid group $Br(M^m, \Sigma_m)$ is called the *braid group on m strings of the manifold M* and is denoted by $Br(m, M)$. Analogously, we call the group $P(M^m, \Sigma_m)$ the *pure braid group on m strings of the manifold M* and denote it by $P(m, M)$. These definitions of braid groups were given by R. Fox and L. Neuwirth [86].

7.2. Artin-Brieskorn braid groups. The braid groups are included in the series of so called generalized braid groups (this was their name in the work of E. Brieskorn of 1971 [42]), or Artin groups (so they were called by E. Brieskorn and K. Saito in the paper of 1972 [45]). They were defined by E. Brieskorn [42], so we call them Artin-Brieskorn groups.

Let V be a finite dimensional real vector space ($\dim V = n$) with Euclidean structure. Let W be a finite subgroup of $GL(V)$ generated by reflections. Let $\mathcal{M}$ be the set of hyperplanes such that W is generated by orthogonal reflections with respect to $M \in \mathcal{M}$. We suppose that for every $w \in W$ and every hyperplane $M \in \mathcal{M}$ the hyperplane $w(M)$ belongs to $\mathcal{M}$.

The group W is generated by the reflections $w_i = w_i(M_i), i \in I$, satisfying only the following relations

$$(w_i w_j)^{m_{i,j}} = e, \quad i, j \in I,$$

where the natural numbers $m_{i,j} = m_{j,i}$ form the *Coxeter matrix* of W from which the *Coxeter graph* $\Gamma(W)$ of W is constructed [41]. We use the following notation of P. Deligne [73]: $\text{prod}(m; x, y)$ denotes the product $xyxy \dots$ (m factors). The *generalized braid group* (or *Artin-Brieskorn group*) $Br(W)$ of W [42], [73] is defined as the group with generators $\{s_i, i \in I\}$ and relations:

$$\text{prod}(m_{i,j}; s_i, s_j) = \text{prod}(m_{j,i}; s_j, s_i).$$

From this we obtain the presentation of the group W by adding the relations:

$$s_i^2 = e; \quad i \in I.$$

We see in Theorem 7.1 that this definition of the generalized braid group agrees with our general definition of a braid group of an action of a group W (Subsection 7.1). We denote by τ_W the canonical homomorphism from $Br(W)$ to W . The classical braids on k strings Br_k are obtained by this construction if W is the symmetric group on $k + 1$ symbols. In this case $m_{i,i+1} = 3$, and $m_{i,j} = 2$ if $j \neq i, i + 1$.

Classification of *irreducible* (with connected Coxeter graph) Coxeter groups is well known (see for example Theorem 1, Chapter VI, §4 of [Bo]). It consists of the three infinite series: A , C (which is also denoted by B because in the corresponding classification of simple Lie algebras two different series B and C have this group as their Weyl group) and D as well as the exceptional groups $E_6, E_7, E_8, F_4, G_2, H_3, H_4$ and $I_2(p)$.

Now let us consider the complexification V_C of the space V and the complexification M_C of $M \in \mathcal{M}$. Let $Y_W = V_C - \bigcup_{M \in \mathcal{M}} M_C$. The group W acts freely on Y_W . Let $X_W = Y_W/W$ then Y_W is a covering over X_W corresponding to the group W . Let $y_0 \in A_0$ be a point in some chamber A_0 and let x_0 stand for its image in X_W . We are in the situation described in Subsection 7.1 in the definition of the braid group of the action of the group W . This braid group is defined as the fundamental group of the space of regular orbits of the action of W . In our case $ORB(V_C, W) = X_W$. So, the generalized braid group is $\pi_1(X_W, x_0)$. For each $j \in I$, let ℓ'_j be the homotopy class of paths in Y_W starting from y_0 and ending in $w_j(y_0)$ which contains a polygon line with successive vertices: $y_0, y_0 + iy_0, w_j(y_0) + iy_0, w_j(y_0)$. The image ℓ_j of the class ℓ'_j in X_W is a loop with base point x_0 .

Theorem 7.1. *The fundamental group $\pi_1(X_W, x_0)$ is generated by the elements ℓ_j satisfying the following relations:*

$$\text{prod}(m_{j,k}; \ell_j, \ell_k) = \text{prod}(m_{k,j}; \ell_k, \ell_j).$$

This theorem was proved by E. Brieskorn [43].

The word problem and the conjugacy problem for Artin-Brieskorn groups were solved by E. Brieskorn and K. Saito [45] and P. Deligne [73]. The biautomatic structure of these groups was established by R. Charney [51].

In the case when V is complex finite dimensional space and W is a finite subgroup of $GL(V)$ generated by *pseudo-reflections* the corresponding braid groups were studied by M. Broué, G. Malle and R. Rouquier [47] and also by D. Bessis and J. Michel [27].

7.3. Braid groups of surfaces. Braid groups of a sphere $Br_n(S^2)$ also have simple geometric interpretation as a group of isotopy classes of braids lying in a layer between two concentric spheres. It has the presentation with generators δ_i , $i = 1, \dots, n-1$, and relations:

$$(7.2) \quad \begin{cases} \delta_i \delta_j &= \delta_j \delta_i, \text{ if } |i-j| > 1, \\ \delta_i \delta_{i+1} \delta_i &= \delta_{i+1} \delta_i \delta_{i+1}, \\ \delta_1 \delta_2 \dots \delta_{n-2} \delta_{n-1}^2 \delta_{n-2} \dots \delta_2 \delta_1 &= 1. \end{cases}$$

This presentation was found by O. Zariski [201] in 1936 and then rediscovered by E. Fadell and J. Van Buskirk [81] in 1961.

Presentations of braid groups of all closed surfaces were obtained by G. P. Scott [181] but look rather complicated.

7.4. Braid groups in handlebodies. The subgroup $Br_{1,n+1}$ of the braid group Br_{n+1} consisting of braids with the fixed first string can be interpreted also as the braid group in a solid torus. Here we study braids in a handlebody of the arbitrary genus g .

Let H_g be a handlebody of genus g . The braid group Br_n^g on n strings in H_g was first considered by A. B. Sossinsky [183]. Let Q_g denote a subset of the complex plane $\mathbb{C}$, consisting of g different points, $Q_g = \{z_1^0, \dots, z_g^0\}$, say, $z_i^0 = i$. The interior of the handlebody H_g may be interpreted as the direct product of the complex plane $\mathbb{C}$ without g points: $\mathbb{C} \setminus Q_g$, and the open interval, for example, $(-1, 1)$:

$$\dot{H}_g = (\mathbb{C} \setminus Q_g) \times (-1, 1).$$

The space $F(\mathbb{C} \setminus Q_g, n)$ can be interpreted as the complement of the arrangement of hyperplanes in $\mathbb{C}^{g+n}$ given by the formulas:

$$H_{j,k} : z_j - z_k = 0 \text{ for all } j, k;$$

$$H_j^i : z_j = z_i^0 \text{ for } i = 1, \dots, g; j = 1, \dots, n.$$

The braids in Br_n^g are considered as lying between the planes with coordinates $z = 0$ and $z = 1$ and connecting the points $((g+1, 0), \dots, (g+n, 0))$. So Br_n^g can be considered as a subgroup of the classical braid group Br_{g+n} on $g+n$ strings such that the braids from Br_n^g leave the first g strings unbraided. In this subsection we denote by $\bar{\sigma}_j$ the standard generators of the group Br_{g+n} . Let τ_k , $k = 1, 2, \dots, g$, be the following braids:

$$\tau_k = \bar{\sigma}_g \bar{\sigma}_{g-1} \dots \bar{\sigma}_{k+1} \bar{\sigma}_k^2 \bar{\sigma}_{k+1}^{-1} \dots \bar{\sigma}_{g-1}^{-1} \bar{\sigma}_g^{-1}.$$

Such a braid is depicted in Figure 7.1. The elements τ_k , $k = 1, 2, \dots, g$, generate a free subgroup F_g in the braid group Br_{g+n} . It follows for example from the Markov normal form that the elements τ_k , $k = 1, 2, \dots, g$, together with the standard generators $\bar{\sigma}_{g+1}, \dots, \bar{\sigma}_{g+n-1}$ generate the group Br_n^g . So, the braid group in the handlebody Br_n^g can be considered as a subgroup of Br_{g+n} , generated by two subgroups: F_g and Br_n . Denote by $\sigma_1, \dots, \sigma_{n-1}$ the standard generators of Br_n considered as the elements of Br_n^g , $\sigma_i = \bar{\sigma}_{g+i}$, $i = 1, \dots, n-1$. So we have




FIGURE 7.1.




FIGURE 7.2.

the presentation of Br_n^g with the generators τ_k and σ_i and relations [183], [190], [193]:

$$(7.3) \quad \begin{cases} \sigma_i \sigma_j &= \sigma_j \sigma_i \text{ if } |i - j| > 1, \\ \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1}, \\ \tau_k \sigma_i &= \sigma_i \tau_k \text{ if } k \geq 1, i \geq 2, \\ \tau_k \sigma_1 \tau_k \sigma_1 &= \sigma_1 \tau_k \sigma_1 \tau_k, \quad k = 1, 2, \dots, g, \\ \tau_k \sigma_1^{-1} \tau_{k+l} \sigma_1 &= \sigma_1^{-1} \tau_{k+l} \sigma_1 \tau_k, \quad k = 1, 2, \dots, g-1; \quad l = 1, 2, \dots, g-k. \end{cases}$$

The relation of the fourth type in (7.3) is the relation of the braid group of type B (C). The relations of the fifth type in (7.3) describe the interaction between the generators of the free group and their closest neighbour σ_1 . Geometrically this is seen in Figure 7.2. If we introduce the new generators θ_k , $k = 1, 2, \dots, g-1$; by the formulas:

$$\theta_k = \sigma_1^{-1} \tau_k \sigma_1$$

we obtain the “positive” presentation of the group B_n^g with generators of the types σ_i , τ_k , θ_k and relations:

$$(7.4) \quad \begin{cases} \sigma_i \sigma_j &= \sigma_j \sigma_i \text{ if } |i - j| > 1, \\ \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1}, \\ \tau_k \sigma_i &= \sigma_i \tau_k \text{ if } k \geq 1, i \geq 2, \\ \tau_k \sigma_1 \tau_k \sigma_1 &= \sigma_1 \tau_k \sigma_1 \tau_k, k = 1, 2, \dots, g, \\ \tau_k \theta_{k+l} &= \theta_{k+l} \tau_k, k = 1, 2, \dots, g-1; l = 1, 2, \dots, g-k. \\ \sigma_1 \theta_k &= \tau_k \sigma_1, k = 1, 2, \dots, g-1. \end{cases}$$

There is a version of Markov Theorem 3.2 for the group Br_n^g [190], [193].

7.5. Braids with singularities. Let BP_n be the subgroup of $\text{Aut } F_n$, generated by both sets of the automorphisms σ_i of (3.6) and ξ_i of the following form:

$$\begin{cases} x_i &\mapsto x_{i+1}, \\ x_{i+1} &\mapsto x_i, \\ x_j &\mapsto x_j, j \neq i, i+1, \end{cases}$$

This is the *braid-permutation group*. R. Fenn, R. Rimányi and C. Rourke proved [83], [84] that this group is given by the set of generators: $\{\xi_i, \sigma_i, i = 1, 2, \dots, n-1\}$ and relations:

$$\begin{cases} \xi_i^2 &= 1, \\ \xi_i \xi_j &= \xi_j \xi_i, \text{ if } |i - j| > 1, \\ \xi_i \xi_{i+1} \xi_i &= \xi_{i+1} \xi_i \xi_{i+1}. \end{cases}$$

The symmetric group relations

$$\begin{cases} \sigma_i \sigma_j &= \sigma_j \sigma_i, \text{ if } |i - j| > 1, \\ \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1}. \end{cases}$$

The braid group relations

$$\begin{cases} \sigma_i \xi_j &= \xi_j \sigma_i, \text{ if } |i - j| > 1, \\ \xi_i \xi_{i+1} \sigma_i &= \sigma_{i+1} \xi_i \xi_{i+1}, \\ \sigma_i \sigma_{i+1} \xi_i &= \xi_{i+1} \sigma_i \sigma_{i+1}. \end{cases}$$

The mixed relations

R. Fenn, R. Rimányi and C. Rourke also gave a geometric interpretation of BP_n as a group of *welded braids*. At first they defined a *welded braid diagram* on n strings as a collection of n monotone arcs starting from n points at a horizontal line of a plane (the top of the diagram) and going down to n points at another horizontal line (the bottom of the diagram). The diagrams may have crossings of two types: 1) the same as usual braids as for example on the Figure 3.2 or 2) welds as depicted in Figure 7.3

Composition of welded braid diagrams on n strings is defined by stacking one diagram under the other. The diagram with no crossings or welds is the identity with respect to composition. So the set of welded braid diagrams on n strings forms a semigroup which is denoted by WD_n .

R. Fenn, R. Rimányi and C. Rourke defined the allowable moves on welded braid diagrams. They consist of usual Reidemeister moves (Figure 3.4) and the specific moves depicted of Figures 7.4, 7.5, 7.6 The automorphisms of F_n which lie in BP_n can be characterized as




FIGURE 7.3.




FIGURE 7.4.




FIGURE 7.5.




FIGURE 7.6.

follows. Let $\pi \in \Sigma_n$ be a permutation and $w_i, i = 1, 2, \dots, n$, be words in F_n . Then the mapping

$$x_i \mapsto w_i^{-1} x_{\pi(i)} w_i$$

determines an injective endomorphism of F_n . If it is also surjective, we call it an automorphism of *permutation-conjugacy type*. The automorphisms of this type comprise a subgroup of $\text{Aut } F_n$ which is precisely BP_n .

The *Baez–Birman monoid* SB_n or *singular braid monoid* [10], [33] is defined as the monoid with generators $g_i, g_i^{-1}, a_i, i = 1, \dots, n-1$, and relations

$$g_i g_j = g_j g_i, \text{ if } |i - j| > 1,$$




FIGURE 7.7.

$$a_i a_j = a_j a_i, \text{ if } |i - j| > 1,$$

$$a_i g_j = g_j a_i, \text{ if } |i - j| \neq 1,$$

$$g_i g_{i+1} g_i = g_{i+1} g_i g_{i+1},$$

$$g_i g_{i+1} a_i = a_{i+1} g_i g_{i+1},$$

$$g_{i+1} g_i a_{i+1} = a_i g_{i+1} g_i,$$

$$g_i g_i^{-1} = g_i^{-1} g_i = 1.$$

In pictures g_i corresponds to canonical generator of the braid group and a_i represents an intersection of the i th and $(i + 1)$ st strand as in Figure 7.7. More detailed geometric interpretation of the Baez–Birman monoid can be found in the article of J. Birman [33]. R. Fenn, E. Keyman and C. Rourke proved [82] that the Baez–Birman monoid embeds in a group SG_n which they called the *singular braid group*:

$$SB_n \rightarrow SG_n.$$

So, in SG_n the elements a_i become invertible and all relations of SB_n remain true.

The analogue of the Birman-Ko-Lee presentation for the singular braid monoid was obtained in [199]. Namely, it was proved that the monoid SB_n has a presentation with generators a_{ts} , a_{ts}^{-1} for $1 \leq s < t \leq n$ and b_{qp} for $1 \leq p < q \leq n$ and relations

$$(7.5) \quad \left\{ \begin{array}{ll} a_{ts} a_{rq} &= a_{rq} a_{ts} \text{ for } (t-r)(t-q)(s-r)(s-q) > 0, \\ a_{ts} a_{sr} &= a_{tr} a_{ts} = a_{sr} a_{tr} \text{ for } 1 \leq r < s < t \leq n, \\ a_{ts} a_{ts}^{-1} &= a_{ts}^{-1} a_{ts} = 1 \text{ for } 1 \leq s < t \leq n, \\ a_{ts} b_{rq} &= b_{rq} a_{ts} \text{ for } (t-r)(t-q)(s-r)(s-q) > 0, \\ a_{ts} b_{ts} &= b_{ts} a_{ts} \text{ for } 1 \leq s < t \leq n, \\ a_{ts} b_{sr} &= b_{tr} a_{ts} \text{ for } 1 \leq r < s < t \leq n, \\ a_{sr} b_{tr} &= b_{ts} a_{sr} \text{ for } 1 \leq r < s < t \leq n, \\ a_{tr} b_{ts} &= b_{sr} a_{tr} \text{ for } 1 \leq r < s < t \leq n, \\ b_{ts} b_{rq} &= b_{rq} b_{ts} \text{ for } (t-r)(t-q)(s-r)(s-q) > 0. \end{array} \right.$$

The elements a_{ts} are defined the same way as in 3.10 and the elements b_{qp} for $1 \leq p < q \leq n$ are defined by

$$(7.6) \quad b_{qp} = (\sigma_{q-1} \sigma_{q-2} \cdots \sigma_{p+1}) x_p (\sigma_{p+1}^{-1} \cdots \sigma_{q-2}^{-1} \sigma_{q-1}^{-1}) \text{ for } 1 \leq p < q \leq n.$$

Geometrically the generators $b_{s,t}$ are depicted in Figure 7.8.




FIGURE 7.8.

8. HOMOLOGICAL PROPERTIES

8.1. Configuration spaces and $K(\pi, 1)$ -spaces. Let $(q_i)_{i \in \mathbb{N}}$ be a fixed sequence of distinct points in the manifold M and put $Q_m = \{q_1, \dots, q_m\}$. We use

$$Q_{m,l} = (q_{l+1}, \dots, q_{l+m}) \in F(M \setminus Q_l, m)$$

as the standard base point of the space $F(M \setminus Q_l, m)$. If $k < m$ we define the projection

$$\text{proj} : F(M \setminus Q_l, m) \rightarrow F(M \setminus Q_l, k)$$

by the formula: $\text{proj}(p_1, \dots, p_m) = (p_1, \dots, p_k)$. The following theorems were proved by E. Fadell and L. Neuwirth [80].

Theorem 8.1. *The triple $\text{proj} : F(M \setminus Q_l, m) \rightarrow F(M \setminus Q_l, k)$ is a locally trivial fiber bundle with fibre $\text{proj}^{-1} Q_{k,l}$ homeomorphic to $F(M \setminus Q_{k+l}, m - k)$.*

Consideration of the sequence of fibrations

$$\begin{aligned} F(M \setminus Q_{m-1}, 1) &\rightarrow F(M \setminus Q_{m-2}, 2) \rightarrow M \setminus Q_{m-2}, \\ F(M \setminus Q_{m-2}, 2) &\rightarrow F(M \setminus Q_{m-3}, 3) \rightarrow M \setminus Q_{m-3}, \\ &\dots, \\ F(M \setminus Q_1, m-1) &\rightarrow F(M, m) \rightarrow M \end{aligned}$$

leads to the following theorem.

Theorem 8.2. *For any manifold M*

$$\pi_i(F(M \setminus Q_1, m-1)) = \oplus_{k=1}^{m-1} \pi_i(M \setminus Q_k)$$

for $i \geq 2$. If $\text{proj} : F(M, m) \rightarrow M$ admits a section then

$$\text{proj}_i \pi_i(F(M, m)) = \oplus_{k=0}^{m-1} \pi_i(M \setminus Q_k), \quad i \geq 2.$$

Corollary 8.1. *If M is the Euclidean r -space, then*

$$\pi_i(F(M, m)) = \oplus_{k=0}^{m-1} \pi_i(\underbrace{S^{r-1} \vee \dots \vee S^{r-1}}_k), \quad i \geq 2.$$

Corollary 8.2. *If M is the Euclidean 2-space, then $F(\mathbb{R}^2, m)$ is the $K(P_m, 1)$ -space and $B(\mathbb{R}^2, m)$ is the $K(Br_m, 1)$ -space.*

Let X_W be the space defined in Subsection 7.2.

Theorem 8.3. *The universal covering of X_W is contractible, and so X_W is a $K(\pi; 1)$ -space.*

This theorem for the groups of types C_n , G_2 and $I_2(p)$, was proved by E. Brieskorn [42] in much the same way as E. Fadell and L. Neuwirth [80] proved Theorems 8.1, 8.2 and Corollary 8.2. For the groups of types D_n and F_4 E. Brieskorn used this method with minor modifications. In general case Theorem 8.3 was proved by P. Deligne [73].

It follows from Theorem 8.2 that $F(\mathbb{C} \setminus Q_g, n)$ and $B(\mathbb{C} \setminus Q_g)$ are $K(\pi, 1)$ -spaces, $\pi_1 B(\mathbb{C} \setminus Q_g) = Br_n^g$, so, $B(\mathbb{C} \setminus Q_g)$ can be considered as the classifying space of Br_n^g .

8.2. Cohomology of pure braid groups. Cohomology of pure braid groups were first calculated by V. I. Arnold [4]. The map

$$\phi : S^{n-1} \rightarrow F(\mathbb{R}^n, 2),$$

described by the formula $\phi(x) = (x, -x)$, is a Σ_2 -equivariant homotopy equivalence. Denote by A the generator of $H^{n-1}(F(\mathbb{R}^n, 2), \mathbb{Z})$ that is mapped by ϕ^* to the standard generator of $H^{n-1}(S^{n-1}, \mathbb{Z})$. For i and j , such that $1 \leq i, j \leq m$, $i \neq j$, specify $\pi_{i,j} : F(\mathbb{R}^n, m) \rightarrow F(\mathbb{R}^n, 2)$ by the formula $\pi_{i,j}(p_1, \dots, p_m) = (p_i, p_j)$. Put

$$A_{i,j} = \pi_{i,j}^*(A) \in H^{n-1}(F(\mathbb{R}^n, m), \mathbb{Z}).$$

It follows that $A_{i,j} = (-1)^n A_{j,i}$ and $A_{i,j}^2 = 0$. For $w \in \Sigma_m$ there is an action $w(A_{i,j}) = A_{w^{-1}(i), w^{-1}(j)}$, since $\pi_{i,j}w = \pi_{w^{-1}(i), w^{-1}(j)}$. Note also that under restriction to

$$F(\mathbb{R}^n \setminus Q_k, m - k) \cong \pi^{-1}(Q_k) \subset F(\mathbb{R}^n, m),$$

the classes $A_{i,j}$ with $1 \leq i, j \leq k$ go to zero since in this case the map $\pi_{i,j}$ is constant on $\pi^{-1}(Q_k)$.

Theorem 8.4. *The cohomology group $H^*(F(\mathbb{R}^n \setminus Q_k, m - k), \mathbb{Z})$ is the free Abelian group with generators*

$$A_{i_1, j_1} A_{i_2, j_2} \dots A_{i_s, j_s},$$

where $k < j_1 < j_2 < \dots < j_s \leq m$ and $i_r < j_r$ for $r = 1, \dots, s$.

The multiplicative structure and the Σ_m -algebra structure of $H^*(F(\mathbb{R}^n, m), \mathbb{Z})$ are given by the following theorem which is proved using the Σ_3 -action on $H^*(F(\mathbb{R}^n, 3), \mathbb{Z})$.

Theorem 8.5. *The cohomology ring $H^*(F(\mathbb{R}^n, m), \mathbb{Z})$ is multiplicatively generated by the square-zero elements*

$$A_{i,j} \in H^{n-1}(F(\mathbb{R}^n, m), \mathbb{Z}), \quad 1 \leq i < j \leq m,$$

subject only to the relations

$$(8.1) \quad A_{i,k} A_{j,k} = A_{i,j} A_{j,k} - A_{i,j} A_{i,k} \text{ for } i < j < k.$$

The Poincaré series for $F(\mathbb{R}^n, m)$ is the product $\prod_{j=1}^{m-1} (1 + jt^{n-1})$.

Remark 8.1. *In the case of $\mathbb{R}^2 = \mathbb{C}$ the cohomology classes $A_{j,k}$ can be interpreted as the classes of cohomology of differential forms*

$$\omega_{j,k} = \frac{1}{2\pi i} \frac{dz_j - dz_k}{z_j - z_k}.$$

E. Brieskorn calculated the cohomology of pure generalized braid groups [42] using ideas of V. I. Arnold for the classical case. Let $\mathcal{V}$ be a finite-dimensional complex vector space and $H_j \in \mathcal{V}$, $j \in I$ be the finite family of complex affine hyperplanes given by linear forms l_j . E. Brieskorn proved the following fact.

Theorem 8.6. *The cohomology classes, corresponding to the holomorphic differential forms*

$$\omega_j = \frac{1}{2\pi i} \frac{dl_j}{l_j},$$

generate the cohomology ring $H^(\mathcal{V} \setminus \cup_{j \in I} H_j, \mathbb{Z})$. Moreover, this ring is isomorphic to the $\mathbb{Z}$ -subalgebra generated by the forms ω_j in the algebra of meromorphic forms on $\mathcal{V}$.*

The cohomologies of pure generalized braid groups are described as follows.

Theorem 8.7. (i) *The cohomology group $H^k(P(W), \mathbb{Z})$ of the pure braid group $P(W)$ with integer coefficients is a free Abelian group, and its rank is equal to the number of elements $w \in W$ of length $l(w) = k$, where l is the length considered with respect to the system of generators consisting of all reflections of W .*

(ii) *The Poincaré series for $H^*(P(W), \mathbb{Z})$ is the product $\prod_{j=1}^n (1 + m_j t)$, where m_j are the exponents of the group W .*

(iii) *The multiplicative structure of $H^*(P(W), \mathbb{Z})$ coincides with the structure of algebra, generated by 1-forms described in the previous theorem.*

8.3. Homology of braid groups. To study the cohomologies of the classical braid groups $H^*(Br_n, \mathbb{Z})$, V. I. Arnold [5] interpreted the space $K(Br_n, 1) \cong B(\mathbb{R}^2, n)$ as the space of monic complex polynomials of degree n without multiple roots

$$P_n(t) = t^n + z_1 t^{n-1} + \dots + z_{n-1} t + z_n.$$

Using this idea he proved Theorems of finiteness, of recurrence and of stabilization. Homology with coefficients in $\mathbb{Z}/2$ were calculated by D. B. Fuks in the following theorems [90].

Theorem 8.8. *The homology of the braid group on the infinite number of strings with coefficients in $\mathbb{Z}/2$ as a Hopf algebra is isomorphic to the polynomial algebra on infinitely many generators $a_i, i = 1, 2, \dots$; $\deg a_i = 2^i - 1$:*

$$H_*(Br_\infty, \mathbb{Z}/2) \cong \mathbb{Z}/2[a_1, a_2, \dots, a_i, \dots]$$

with the coproduct given by the formula:

$$\Delta(a_i) = 1 \otimes a_i + a_i \otimes 1.$$

Theorem 8.9. *The canonical inclusion $Br_n \rightarrow Br_\infty$ induces a monomorphism in homology with coefficients in $\mathbb{Z}/2$. Its image is the subcoalgebra of the polynomial algebra $\mathbb{Z}/2[a_1, a_2, \dots, a_i, \dots]$ with $\mathbb{Z}/2$ -basis consisting of monomials*

$$a_1^{k_1} \dots a_i^{k_i} \text{ such that } \sum_i k_i 2^i \leq n.$$

Theorem 8.10. *The canonical homomorphism $Br_n \rightarrow BO_n, 1 \leq n \leq \infty$ induces a monomorphism (of Hopf algebras if $n = \infty$)*

$$H_*(Br_n, \mathbb{Z}/2) \rightarrow H_*(BO_n, \mathbb{Z}/2).$$

F. R. Cohen calculated the homology of braid groups with coefficients $\mathbb{Z}/p, p > 2$ also as modules over the Steenrod algebra [54], [55], [56].

Later V. V. Goryunov [108], [109] applied the methods of Fuks and expressed the cohomologies of the generalized braid groups of types C and D in terms of the cohomologies of the classical braid groups.




FIGURE 9.1.

9. CONNECTIONS WITH THE OTHER DOMAINS

9.1. Markov Theorem. Suppose a braid depicted in Figure 3.1 is placed in a cube. On the boundary of the cube join the point A_i to the point B_i by a mutually disjoint simple arc D_i . Since our initial braid does not intersect the boundary of the cube except at the points $A_1, \dots, A_n$ and $B_1, \dots, B_n$ we obtain a link (or, in particular, a knot) i.e. a system of simple closed curves in $\mathbb{R}^3$. A link obtained in such a manner is called the *closure* of the braid, see Figure 9.1.

Theorem 9.1. (J. W. Alexander) *Any link can be represented by a closed braid.*

The next step is to understand equivalence classes of braids which correspond to links. The following Markov Theorem gives an answer to this question. At first we define two types of Markov moves for braids.

Type 1 Markov move replaces a braid β on n strings by its conjugate $\gamma\beta\gamma^{-1}$.

Type 2 Markov move replaces a braid β on n strings by the braid $j_n(\beta)\sigma_n$ on $n+1$ strings or by $j_n(\beta)\sigma_n^{-1}$ where j_n is the canonical inclusion of the group Br_n into the group Br_{n+1} (see Subsection 3.1)

$$j_n : Br_n \rightarrow Br_{n+1}.$$

Theorem 9.2. (A. A. Markov) *Suppose that β and β' are two braids (not necessary with the same number of strings). Then, the closures of β and β' represent the same link if and only if β can be transformed into β' by means of a finite number of type 1 and type 2 Markov moves. Namely there exists the following sequence,*

$$\beta = \beta_0 \rightarrow \beta_1 \rightarrow \dots \rightarrow \beta_m = \beta',$$

such that, for $i = 0, 1, \dots, m-1$, β_{i+1} is obtained from β_i by the application of a type 1 or 2 Markov moves or their inverses.

In other words, if we consider the disjoint union of all braid groups

$$\coprod_{n=1}^{n=\infty} Br_n,$$

then the Markov moves of types 1 and 2 define the equivalence relation on this set $\sim$ such that the quotient set

$$\prod_{n=1}^{n=\infty} Br_n / \sim$$

is in one-to-one correspondence with isotopy classes of links.

There exist a lot of proofs of Markov Theorem, see for example the work of P. Traczyk [186].

9.2. Homotopy groups of spheres and Makanin braids. Consider the coordinate projections for the spaces $F(M, m)$ where M is a manifold (see Subsection 7.1)

$$d_i : F(M, n+1) \rightarrow F(M, n), \quad i = 0, \dots, n,$$

defined by the formula

$$d_i(p_1, \dots, p_{i+1}, \dots, p_{n+1}) = (p_1, \dots, \hat{p}_{i+1}, \dots, p_{n+1}).$$

By taking the fundamental group the maps d_i induces group homomorphisms

$$d_{i*} : P_{m+1}(M) \rightarrow P_m(M), \quad i = 0, \dots, n.$$

A braid $\beta \in Br_{n+1}$ is called *Makanin* (*smooth* in the terminology of D. L. Johnson [120], *Brunnian* in the terminology of J. A. Berrick, F. R. Cohen, Y. L. Wong and J. Wu [22]) if $d_i(\beta) = 1$ for all $0 \leq i \leq n$. We call them Makanin, because up to our knowledge it was G. S. Makanin who first mentioned them [127], page 78, question 6.23. In other words the group of Makanin braids $Mak_{n+1}(M)$ is given by the formula

$$Mak_{n+1}(M) = \cap_{i=0}^n \text{Ker}(d_{i*} : P_{m+1}(M) \rightarrow P_m(M)).$$

The canonical embedding of the open disc D^2 into the sphere S^2

$$f : D^2 \rightarrow S^2$$

induces a group homomorphism

$$f_* : Mak_n(D^2) \rightarrow Mak_n(S^2)$$

where $Mak_n(D^2)$ is the Makanin subgroup Mak_n of the classical braid group Br_n . The group Mak_n is free [112, 120]. The following theorem is proved in [22].

Theorem 9.3. *The is an exact sequence of groups*

$$1 \rightarrow Mak_{n+1}(S^2) \rightarrow Mak_n(D^2) \rightarrow Mak_n(S^2) \rightarrow \pi_{n-1}(S^2) \rightarrow 1$$

for $n \geq 5$.

Here as usual $\pi_k(S^2)$ denote the k -th homotopy group of the sphere S^2 .

For instance, $Mak_5(S^2)$ modulo Mak_5 is $\pi_4(S^2) = \mathbb{Z}/2$. The other homotopy groups of S^2 are as follows

$$\pi_5(S^2) = \mathbb{Z}/2, \quad \pi_6(S^2) = \mathbb{Z}/12, \quad \pi_7(S^2) = \mathbb{Z}/2, \quad \pi_8(S^2) = \mathbb{Z}/2, \dots$$

Thus, up to certain range, $Mak_n(S^2)$ modulo Mak_n are known by nontrivial calculation of $\pi_*(S^2)$.

REFERENCES

- [1] *S. I. Adyan*, Fragments of the word Δ in the braid group. (Russian) *Mat. Zametki* 36 (1984), no. 1, 25–34.
- [2] *A. Adem; D. Cohen; F. R. Cohen*, On representations and K -theory of the braid groups. *Math. Ann.* 326 (2003), no. 3, 515–542.
- [3] *J. W. Alexander*, A lemma on systems of knotted curves. *Proc. Nat. Acad. Sci. USA*, 9, 93–95.
- [4] *V. I. Arnold*, The cohomology ring of coloured braids, *Mat. Zametki*, 1969, 5, 227–231 (Russian). English transl. in *Math. Notes*, 1969, 5, 138–140.
- [5] *V. I. Arnold*, On some topological invariants of algebraic functions, *Trudy Moscov. Mat. Obshch.* 1970, 21, 27–46 (Russian), English transl. in *Trans. Moscow Math. Soc.*, 1970, v. 21, 30–52.
- [6] *V. I. Arnold*, Topological invariants of algebraic functions II, *Functional Anal. Appl.* 1970, v. 4, No 2, 1–9 (Russian).
- [7] *E. Artin*, Theorie der Zöpfe, *Abh. Math. Semin. Univ. Hamburg*, 1925, v. 4, 47–72.
- [8] *E. Artin*, Theory of braids, *Ann. Math.* 1947, v. 48, No 1, 101–126.
- [9] *E. Artin*, Braids and permutations. *Ann. of Math.* (2) 48, (1947). 643–649.
- [10] *J. C. Baez*, Link invariants of finite type and perturbation theory. *Lett. Math. Phys.* 26 (1992), no. 1, 43–51.
- [11] *V. G. Bardakov*, On the theory of braid groups. (Russian) *Mat. Sb.* 183 (1992), no. 6, 3–42; translation in *Russian Acad. Sci. Sb. Math.* 76 (1993), no. 1, 123–153.
- [12] *V. G. Bardakov*, The width of verbal subgroups of some Artin groups. (Russian) *Group and metric properties of mappings* (Russian), 8–18, Novosibirsk. Gos. Univ., Novosibirsk, 1995.
- [13] *V. G. Bardakov*, The structure of a group of conjugating automorphisms. (Russian) *Algebra Logika* 42 (2003), no. 5, 515–541, 636; translation in *Algebra Logic* 42 (2003), no. 5, 287–303.
- [14] *V. G. Bardakov*, The virtual and universal braids. *Fund. Math.* 184 (2004), 1–18.
- [15] *V. G. Bardakov*, Linear representations of the braid groups of some manifolds. *Acta Appl. Math.* 85 (2005), no. 1–3, 41–48.
- [16] *V. G. Bardakov*, Linear representations of the group of conjugating automorphisms and the braid groups of some manifolds. (Russian) *Sibirsk. Mat. Zh.* 46 (2005), no. 1, 17–31; translation in *Siberian Math. J.* 46 (2005), no. 1, 13–23
- [17] *P. Bellingeri*, Tresses sur surface et invariants d’entrelacs, Thèse, Univ. Grenoble I (2003).
- [18] *P. Bellingeri*, Centralisers in surface braid groups. *Comm. Algebra* 32 (2004), no. 10, 4099–4115.
- [19] *P. Bellingeri*, On presentations of surface braid groups. *J. Algebra* 274 (2004), no. 2, 543–563.
- [20] *P. Bellingeri; L. Funar*, Braids on surfaces and finite type invariants. *C. R. Math. Acad. Sci. Paris* 338 (2004), no. 2, 157–162.
- [21] *P. Bellingeri, V. Vershinin* Presentations of surface braid groups by graphs, to appear in *Fund. Math.*
- [22] *J. A. Berrick, F. R. Cohen, Y. L. Wong and J. Wu*, Configurations, b braids, and homotopy groups, *JAMS*, to appear.
- [23] *D. Bessis*, David Groupes des tresses et lments rguliers. (French) [Braid groups and regular elements] *J. Reine Angew. Math.* 518 (2000), 1–40.
- [24] *D. Bessis*, Zariski theorems and diagrams for braid groups. *Invent. Math.* 145 (2001), no. 3, 487–507.
- [25] *D. Bessis*, The dual braid monoid. *Ann. Sci. cole Norm. Sup.* (4) 36 (2003), no. 5, 647–683.
- [26] *D. Bessis; F. Digne; J. Michel*, Springer theory in braid groups and the Birman-Ko-Lee monoid. *Pacific J. Math.* 205 (2002), no. 2, 287–309.
- [27] *D. Bessis; J. Michel*, Explicit presentations for exceptional braid groups. *Experiment. Math.* 13 (2004), no. 3, 257–266.
- [28] *S. Bigelow*, The Burau representation is not faithful for $n = 5$. *Geom. Topol.* 3 (1999), 397–404 (electronic).
- [29] *S. Bigelow*, Braid groups are linear, *J. Amer. Math. Soc.* 14 (2001), no. 2, 471–486.
- [30] *J. S. Birman*, On braid groups. *Comm. Pure Appl. Math.* 22 1968 41–72.
- [31] *J. S. Birman*, Mapping class groups and their relationship to braid groups. *Comm. Pure Appl. Math.* 22 1969 213–238.

- [32] *J. S. Birman*, Braids, Links, and Mapping Class Groups, *Ann. Math. Stud.*, No 82, 1974.
- [33] *J. S. Birman*, New points of view in knot theory, *Bull. Amer. Math. Soc.* 1993, 28, No 2, 253–387.
- [34] *J. S. Birman and T. E. Brendle*, BRAIDS: A SURVEY. *Arxiv.math.GT/0409205*
- [35] *J. S. Birman; K. H. Ko; S. J. Lee*, A new approach to the word and conjugacy problems in the braid groups. *Adv. Math.* 139 (1998), no. 2, 322–353.
- [36] *J. S. Birman; K. H. Ko; S. J. Lee*, The infimum, supremum, and geodesic length of a braid conjugacy class. *Adv. Math.* 164 (2001), no. 1, 41–56.
- [37] *J. S. Birman; D. D. Long; J. A. Moody*, Finite-dimensional representations of Artin’s braid group. The mathematical legacy of Wilhelm Magnus: groups, geometry and special functions (Brooklyn, NY, 1992), 123–132, *Contemp. Math.*, 169, Amer. Math. Soc., Providence, RI, 1994.
- [38] *J. S. Birman; W. W. Menasco*, On Markov’s theorem. *Knots 2000 Korea*, Vol. 1 (Yongpyong). *J. Knot Theory Ramifications* 11 (2002), no. 3, 295–310.
- [39] *J. S. Birman and B. Wajnryb*, Markov classes in certain finite quotients of Artin’s braid group. *Israel J. Math.* 56 (1986), no. 2, 160–178.
- [40] *J. S. Birman and H. Wenzl*, Braids, link polynomials and a new algebra. *Trans. Amer. Math. Soc.* 313 (1989), no. 1, 249–273.
- [41] *N. Bourbaki*, Groupes et algèbres de Lie, Chaps. 4–6, Masson, Paris, 1981.
- [42] *E. Brieskorn*, Sur les groupes de tresses [d’après V. I. Arnol’d]. (French) *Séminaire Bourbaki*, 24ème ann(1971/1972), Exp. No. 401, pp. 21–44. *Lecture Notes in Math.*, Vol. 317, Springer, Berlin, 1973.
- [43] *E. Brieskorn*, Die Fundamentalgruppe des Raumes der regulären Orbits einer endlichen komplexen Spiegelungsgruppe. (German) *Invent. Math.* 12 (1971), 57–61.
- [44] *E. Brieskorn*, Automorphic sets and braids and singularities. *Braids* (Santa Cruz, CA, 1986), 45–115, *Contemp. Math.*, 78, Amer. Math. Soc., Providence, RI, 1988.
- [45] *E. Brieskorn and K. Saito*, Artin-Gruppen und Coxeter-Gruppen. (German) *Invent. Math.* 17 (1972), 245–271.
- [46] *C. Broto; V. V. Vershinin*, On the generalized homology of Artin groups. *J. Math. Sci. (N. Y.)* 113 (2003), no. 4, 545–547, translation from Russian: *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 266 (2000), *Teor. Predst. Din. Sist. Komb. i Algoritm. Metody.* 5, 7–12, 336.
- [47] *M. Broué, G. Malle, R. Rouquier* Complex reflection groups, braid groups, Hecke algebras. *J. Reine Angew. Math.* 500, 127–190 (1998).
- [48] *W. Burau*, Über Zopfvarianten. (German) *Abh. Math. Semin. Hamb. Univ.* 9, 117–124 (1932).
- [49] *W. Burau*, Über Verkettungsgruppen. (German) *Abh. Math. Semin. Hamb. Univ.* 11, 171–178 (1935).
- [50] *W. Burau*, Über Zopfgruppen und gleichsinnig verdrehte Verkettungen. (German) *Abh. Math. Semin. Hamb. Univ.* 11, 179–186 (1935).
- [51] *R. Charney*, Artin groups of finite type are biautomatic. *Math. Ann.* 292 (1992), no. 4, 671–683.
- [52] *R. Charney*, Geodesic automation and growth functions for Artin groups of finite type. *Math. Ann.* 301 (1995), no. 2, 307–324.
- [53] *W.-L. Chow*, On the algebraical braid group, *Ann. Math.* 1948, 49, No 3, 654–658.
- [54] *F. Cohen*, Cohomology of braid spaces *Bull. Amer. Math. Soc.* 1973, 79, No 4 763–766.
- [55] *F. Cohen*, Homology of $\Omega^{n+1}\Sigma^{n+1}X$ and $C_{n+1}X$, $n > 0$ *Bull. Amer. Math. Soc.* 1973, 79, No 6, 1236–1241.
- [56] *F. Cohen*, Braid orientations and bundles with flat connections *Invent. Math.* 1978, 46, 99–110.
- [57] *F. Cohen*, Artin’s braid groups, classical homotopy theory, and other curiosities, *Braids* (*Contemp. Math.* **78**, 1988), 167–206.
- [58] *F. Cohen*, On braid groups, homotopy groups, and modular forms. *Advances in topological quantum field theory*, 275–288, *NATO Sci. Ser. II Math. Phys. Chem.*, 179, Kluwer Acad. Publ., Dordrecht, 2004.
- [59] *F. Cohen; T. Lada; J. P. May*, The homology of iterated loop spaces. *Lecture Notes in Mathematics*, Vol. 533. Springer-Verlag, Berlin-New York, 1976. vii+490 pp.
- [60] *F. Cohen; V. V. Vershinin*, Thom spectra which are wedges of Eilenberg-MacLane spectra. *Stable and unstable homotopy* (Toronto, ON, 1996), 43–65, *Fields Inst. Commun.*, 19, Amer. Math. Soc., Providence, RI, 1998.

- [61] *F. Cohen; J. Wu*, On braid groups, free groups, and the loop space of the 2-sphere. Categorical decomposition techniques in algebraic topology (Isle of Skye, 2001), 93–105, Progr. Math., 215, Birkhuser, Basel, 2004.
- [62] *R. Corran*, A normal form for a class of monoids including the singular braid monoids. J. Algebra. 223 (2000), no. 1, 256–282.
- [63] *H. S. M. Coxeter*, Factor groups of the braid group. Proc. Fourth Canad. Math. Congr. Banff, 1957, (1959) 95–122.
- [64] *H. S. M. Coxeter, W. O. J. Moser* Generators and relations for discrete groups. 3rd ed. Ergebnisse der Mathematik und ihrer Grenzgebiete. Band 14. Berlin-Heidelberg-New York: Springer-Verlag. IX, 161 p. (1972).
- [65] *J. Crisp; Paris, Luis* The solution to a conjecture of Tits on the subgroup generated by the squares of the generators of an Artin group. Invent. Math. 145 (2001), no. 1, 19–36.
- [66] *O. Dasbach and B. Gemein*, A faithful representation of the singular braid monoid on three strands. Knots in Hellas '98 (Delphi), 48–58, Ser. Knots Everything, 24, World Sci. Publishing, River Edge, NJ, 2000.
- [67] *P. Dehornoy*, Deux propriétés des groupes de tresses. (French) [Two properties of braid groups] C. R. Acad. Sci. Paris S I Math. 315 (1992), no. 6, 633–638.
- [68] *P. Dehornoy*, Braid groups and left distributive operations. Trans. Amer. Math. Soc. 345 (1994), no. 1, 115–150.
- [69] *P. Dehornoy*, Braids and self-distributivity. Progress in Mathematics, 192. Birkhäuser Verlag, Basel, 2000. xx+623 pp.
- [70] *P. Dehornoy; I. Dynnikov; D. Rolfsen; B. Wiest*, Why are braids orderable? Panoramas et Synthèses [Panoramas and Syntheses], 14. Société Mathématique de France, Paris, 2002. xiv+190 pp.
- [71] *P. Dehornoy; Y. Lafont*, Homology of Gaussian groups. Ann. Inst. Fourier (Grenoble) 53 (2003), no. 2, 489–540.
- [72] *P. Dehornoy; L. Paris*, Gaussian groups and Garside groups, two generalisations of Artin groups. Proc. London Math. Soc. (3) 79 (1999), no. 3, 569–604.
- [73] *P. Deligne*, Les immeubles des groupes de tresses généralisés. (French) Invent. Math. 17 (1972), 273–302.
- [74] *G. Dethloff; S. Orevkov; M. Zaidenberg*, Plane curves with a big fundamental group of the complement. Voronezh Winter Mathematical Schools, 63–84, Amer. Math. Soc. Transl. Ser. 2, 184, Amer. Math. Soc., Providence, RI, 1998.
- [75] *J. Daz-Cantos; J. Gonzalez-Meneses; J. M. Tornero*, On the singular braid monoid of an orientable surface. Proc. Amer. Math. Soc. 132 (2004), no. 10, 2867–2873.
- [76] *E. El-Rifai and H. R. Morton*, Algorithms for positive braids. Quart. J. Math. Oxford Ser. (2) 45 (1994), no. 180, 479–497.
- [77] *M. Eppler*, Orbits of asteroids, a braid, and the first link invariant, Math. Intell. 1998, 20, No 1, 45–52.
- [78] *D. B. A. Epstein; J. W. Cannon; D. E. Holt; S. V. F. Levy; M. S. Paterson; W. P. Thurston*, Word processing in groups. Jones and Bartlett Publishers, Boston, MA, 1992. xii+330 pp.
- [79] *E. Fadell; S. Y. Hussein*, Geometry and topology of configuration spaces. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2001. xvi+313 pp.
- [80] *E. Fadell and L. Neuwirth*, Configuration spaces, Math. Scand. 1962, 10, Fasc. I, 111–118.
- [81] *E. Fadell and J. Van Buskirk*, The braid groups of E^2 and S^2 . Duke Math. J. 29 1962, 243–257.
- [82] *R. Fenn; E. Keyman; C. Rourke*, The singular braid monoid embeds in a group. J. Knot Theory Ramifications 7 (1998), no. 7, 881–892.
- [83] *R. Fenn; R. Rimányi; C. Rourke*, Some remarks on the Braid-Permutation Group. Topics in Knot Theory. Kluwer Academic Publishers, 1993, 57–68.
- [84] *R. Fenn; R. Rimányi; C. Rourke*, The braid-permutation group. Topology 36 (1997), no. 1, 123–135.
- [85] *R. Fenn; D. Rolfsen; J. Zhu*, Centralisers in the braid group and singular braid monoid. Enseign. Math. (2) 42 (1996), no. 1-2, 75–96.
- [86] *R. Fox and L. Neuwirth*, The braid groups, Math. Scand. 1962, 10, Fasc. I, 119–126.

- [87] *N. Franco; J. González-Meneses*, Computation of centralizers in braid groups and Garside groups. Proceedings of the International Conference on Algebraic Geometry and Singularities (Spanish) (Sevilla, 2001). Rev. Mat. Iberoamericana 19 (2003), no. 2, 367–384.
- [88] *N. Franco; J. González-Meneses*, Conjugacy problem for braid groups and Garside groups. J. Algebra 266 (2003), no. 1, 112–132.
- [89] *R. Fricke, F. Klein*, Vorlesungen über die Theorie der Automorphen Functionen. Bd. I. Gruppentheoretischen Grundlagen. Teubner, Leipzig, 1897 (Johnson Repr. Corp., N. Y., 1965).
- [90] *D. B. Fuks*, Cohomology of the braid group mod 2, Functional. Anal. i Prilozh. 1970, 4, No 2, 62–75 (Russian), English transl. in Functional Anal. Appl. 4 (1970), 143–151.
- [91] *F. A. Garside*, The braid group and other groups, Quart. J. Math. Oxford Ser. 1969, 20, 235–254.
- [92] *B. J. Gassner*, On braid groups. Abh. Math. Sem. Univ. Hamburg, 25, 1961, 10–22.
- [93] *B. Gemein*, Singular braids and Markov’s theorem. J. Knot Theory Ramifications 6 (1997), no. 4, 441–454.
- [94] *B. Gemein*, Representations of the singular braid monoid and group invariants of singular knots. Topology Appl. 114 (2001), no. 2, 117–140.
- [95] *E. Godelle; L. Paris*, On singular Artin monoids. Geometric methods in group theory, 43–57, Contemp. Math., 372, Amer. Math. Soc., Providence, RI, 2005.
- [96] *V. A. Golubeva; V. P. Leksin*, Quadratic relations in the cohomology of generalized colored braids, and Dunkl identities. (Russian) Funktsional. Anal. i Prilozhen. 30 (1996), no. 2, 73–76; translation in Funct. Anal. Appl. 30 (1996), no. 2, 131–134.
- [97] *V. A. Golubeva; V. P. Leksin*, On two types of representations of the braid group associated with the Knizhnik-Zamolodchikov equation of the B_n type. J. Dynam. Control Systems 5 (1999), no. 4, 565–596.
- [98] *D. L. Gonçalves; J. Guaschi*, On the structure of surface pure braid groups. J. Pure Appl. Algebra 182 (2003), no. 1, 33–64.
- [99] *D. L. Gonçalves; J. Guaschi*, The roots of the full twist for surface braid groups. Math. Proc. Cambridge Philos. Soc. 137 (2004), no. 2, 307–320.
- [100] *D. L. Gonçalves; J. Guaschi*, The braid groups of the projective plane. Algebr. Geom. Topol. 4 (2004), 757–780 (electronic).
- [101] *D. L. Gonçalves; J. Guaschi*, The braid group $B_{n,m}(\mathbb{S}^2)$ and a generalisation of the Fadell-Neuwirth short exact sequence. J. Knot Theory Ramifications 14 (2005), no. 3, 375–403.
- [102] *J. González-Meneses*, New presentations of surface braid groups. J. Knot Theory Ramifications 10 (2001), no. 3, 431–451.
- [103] *J. González-Meneses*, Ordering pure braid groups on compact, connected surfaces. Pacific J. Math. 203 (2002), no. 2, 369–378.
- [104] *J. González-Meneses*, Presentations for the monoids of singular braids on closed surfaces. Comm. Algebra 30 (2002), no. 6, 2829–2836.
- [105] *J. González-Meneses*, The n th root of a braid is unique up to conjugacy. Algebr. Geom. Topol. 3 (2003), 1103–1118.
- [106] *J. González-Meneses*, Improving an algorithm to solve multiple simultaneous conjugacy problems in braid groups. Geometric methods in group theory, 35–42, Contemp. Math., 372, Amer. Math. Soc., Providence, RI, 2005.
- [107] *J. González-Meneses; L. Paris*, Vassiliev invariants for braids on surfaces. Trans. Amer. Math. Soc. 356 (2004), no. 1, 219–243.
- [108] *V. V. Goryunov*, The cohomology of braid groups of series C and D and certain stratifications. (Russian) Funktsional. Anal. i Prilozhen. 12 (1978), no. 2, 76–77.
- [109] *V. V. Goryunov*, Cohomology of braid groups of series C and D . (Russian) Trudy Moskov. Mat. Obshch. 42 (1981), 234–242.
- [110] *J. Guaschi*, Representations of Artin’s braid groups and linking numbers of periodic orbits. J. Knot Theory Ramifications 4 (1995), no. 2, 197–212.
- [111] *J. Guaschi*, Nielsen theory, braids and fixed points of surface homeomorphisms. Topology Appl. 117 (2002), no. 2, 199–230.
- [112] *G. G. Gurzo*, On the group of smooth braids. (Russian) 16th All-Union Algebra Conference, Abstracts II, p. 39–40. Leningrad, 1981.

- [113] *G. G. Gurzo*, Systems of generators for normalizers of certain elements of a braid group. (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* 48 (1984), no. 3, 476–519.
- [114] *G. G. Gurzo*, Centralizers of finite sets of elements of the braid group $\mathfrak{B}_{n+1}$. (Russian) *Mat. Zametki* 37 (1985), no. 1, 3–6, 137.
- [115] *G. G. Gurzo*, Systems of generators for the centralizers of the rigid elements of a braid group. (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* 51 (1987), no. 5, 915–935,
- [116] *V. Hansen*, Braids and coverings: selected topics. London Mathematical Society Student Text, 18. Cambridge etc.: Cambridge University Press. x, 191 p. (1989).
- [117] *A. Hurwitz*, Über Riemannische Flächen mit gegebenen Verzweigungspunkten, *Math. Ann.* 1891, 39, 1–61.
- [118] *N. V. Ivanov*, Algebraic properties of the Teichmüller modular group. (Russian) *Dokl. Akad. Nauk SSSR* 275 (1984), no. 4, 786–789.
- [119] *A. J  rai Jr.*, On the monoid of singular braids. *Topology Appl.* 96 (1999), no. 2, 109–119.
- [120] *D. L. Johnson*, Towards a characterization of smooth braids. *Math. Proc. Cambridge Philos. Soc.* 92 (1982), no. 3, 425–427.
- [121] *V. F. R. Jones*, Hecke algebra representations of braid groups and link polynomials, *Ann. Math.* 1987, 126, 335–388.
- [122] *A. Joyal; R. Street*, Braided tensor categories. *Adv. Math.* 102 (1993), no. 1, 20–78.
- [123] *S. Kamada*, Braid presentation of virtual knots and welded knots. *Arxiv.math.GT/0008092*.
- [124] *S. Kamada; Y. Matsumoto*, Certain racks associated with the braid groups. *Knots in Hellas '98 (Delphi)*, 118–130, *Ser. Knots Everything*, 24, World Sci. Publishing, River Edge, NJ, 2000.
- [125] *L. H. Kauffman; S. Lambropoulou*, Virtual braids. *Fund. Math.* 184 (2004), 159–186.
- [126] *F. Klein*, Vorlesungen   ber h  here Geometrie. 3. Aufl., bearbeitet und herausgegeben von *W. Blaschke*. (German) VIII + 405 S. Berlin, J. Springer (Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen Bd. 22) (1926).
- [127] Kourouva notebook: unsolved problems in group theory. Seventh edition. 1980. *Akad. Nauk SSSR Sibirsk. Otdel., Inst. Mat., Novosibirsk*, 1980. 115 pp. (Russian)
- [128] *D. Kramer*, The braid group B_4 is linear, *Invent. Math.* 142 (2000), no. 3, 451–486.
- [129] *D. Kramer*, Braid groups are linear, *Ann. of Math.* (2) 155 (2002), no. 1, 131–156.
- [130] *Vik. S. Kulikov*, A full-twist factorization formula with a double number of strings. (Russian) *Izv. Ross. Akad. Nauk Ser. Mat.* 68 (2004), no. 1, 123–158; translation in *Izv. Math.* 68 (2004), no. 1, 125–158.
- [131] *Vik. S. Kulikov; V. M. Kharlamov*, On braid monodromy factorizations. (Russian) *Izv. Ross. Akad. Nauk Ser. Mat.* 67 (2003), no. 3, 79–118; translation in *Izv. Math.* 67 (2003), no. 3, 499–534.
- [132] *S. Lambropoulou*, Solid torus links and Hecke algebras of $\mathcal{B}$ -type. *Proceedings of the Conference on Quantum Topology (Manhattan, KS, 1993)*, 225–245, World Sci. Publishing, River Edge, NJ, 1994.
- [133] *S. Lambropoulou; C. P. Rourke*, Markov’s theorem in 3-manifolds. Special issue on braid groups and related topics (Jerusalem, 1995). *Topology Appl.* 78 (1997), no. 1-2, 95–122.
- [134] *R. J. Lawrence*, Homological representations of the Hecke algebra. *Comm. Math. Phys.* 135 (1990), no. 1, 141–191.
- [135] *V. Ya. Lin*, Artinian braids and groups and spaces connected with them, *Itogi Nauki i Tekhniki (Algebra, Topologiya, Geometriya)* 1979, 17, 159–227 (Russian). English transl. in *J. Soviet Math.* 18 (1982) 736–788.
- [136] *V. Ya. Lin*, Configuration spaces of $\mathbb{C}$ and $\mathbb{C}P^1$: some analytic properties, *Arxiv: math.AG/0403120*
- [137] *V. Ya. Lin*, Braids and Permutations, *Arxiv: math.GR/0404528*
- [138] *D. D. Long and M. Paton*, The Burau representation is not faithful for $n \geq 6$. *Topology* 32 (1993), no. 2, 439–447.
- [139] *W. Magnus*,   ber Automorphismen von Fundamentalgruppen berandeter Fl  chen. (German) *Math. Ann.* 109, 617–646 (1934).
- [140] *W. Magnus* Braid groups: a survey. *Proc. 2nd internat. Conf. Theory of Groups, Canberra 1973*, *Lect. Notes Math.* 372, 463–487 (1974)

- [141] *W. Magnus; A. Karrass; D. Solitar*, Combinatorial group theory. Presentations of groups in terms of generators and relations. 2nd rev. ed. Dover Books on Advanced Mathematics. New York: Dover Publications, Inc. XII, 444 p. (1976).
- [142] *W. Magnus and A. Peluso*, On knot groups. *Comm. Pure Appl. Math.* 20, 1967, 749–770.
- [143] *G. S. Makanin*, The conjugacy problem in the braid group, (Russian) *Dokl. Akad. Nauk SSSR*. 182, (1968) 495–496.
- [144] *G. S. Makanin*, The normalizers of a braid group. (Russian) *Mat. Sb. (N.S.)* 86(128) (1971), 171–179.
- [145] *G. S. Makanin*, Separable closed braids. (Russian) *Mat. Sb. (N.S.)* 132(174) (1987), no. 4, 531–540; translation in *Math. USSR-Sb.* 60 (1988), no. 2, 521–531.
- [146] *G. S. Makanin*, A representation of an oriented knot. (Russian) *Dokl. Akad. Nauk SSSR* 299 (1988), no. 5, 1060–1063; translation in *Soviet Math. Dokl.* 37 (1988), no. 2, 522–525.
- [147] *G. S. Makanin*, An analogue of the Alexander-Markov theorem. (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* 53 (1989), no. 1, 200–210; translation in *Math. USSR-Izv.* 34 (1990), no. 1, 201–211.
- [148] *G. S. Makanin; A. G. Savushkina*, An equation in a free group that defines colored braids. (Russian) *Mat. Zametki* 70 (2001), no. 4, 591–602; translation in *Math. Notes* 70 (2001), no. 3-4, 535–544
- [149] *S. Manfardini*, Some subgroups of Artin’s braid group. Special issue on braid groups and related topics (Jerusalem, 1995). *Topology Appl.* 78 (1997), no. 1-2, 123–142.
- [150] *A. A. Markoff*, Über die freie Äquivalenz der geschlossenen Zöpfe, *Math. Sb.* 1936, 16, 73–78.
- [151] *A. A. Markoff*, Foundations of the Algebraic Theory of Tresses, *Trudy Mat. Inst. Steklova*, No 16, 1945 (Russian, English summary).
- [152] *J. McCarthy*, A "Tits-alternative" for subgroups of surface mapping class groups. *Trans. Amer. Math. Soc.* 291 (1985), no. 2, 583–612.
- [153] *J. Michel*, A note on words in braid monoids. *J. Algebra* 215 (1999), no. 1, 366–377.
- [154] *J. A. Moody*, The Burau representation of the braid group B_n is unfaithful for large n . *Bull. Amer. Math. Soc. (N.S.)* 25 (1991), no. 2, 379–384.
- [155] *S. Moran*, The mathematical theory of knots and braids. An introduction. North-Holland Mathematics Studies, 82. North-Holland Publishing Co., Amsterdam, 1983. xii+295 pp.
- [156] *J. Murakami*, The Kauffman polynomial of links and representation theory. *Osaka J. Math.* 24 (1987), no. 4, 745–758.
- [157] *K. Murasugi; B. Kurpita*, A study of braids. *Mathematics and its Applications*, 484. Kluwer Academic Publishers, Dordrecht, 1999. x+272 pp.
- [158] *F. Napolitano*, Configuration spaces on surfaces. *C. R. Acad. Sci. Paris Sr. I Math.* 327 (1998), no. 10, 887–892.
- [159] *F. Napolitano*, On the cohomology of configuration spaces on surfaces. *J. London Math. Soc. (2)* 68 (2003), no. 2, 477–492.
- [160] *F. Napolitano*, Cohomology rings of spaces of generic bipolynomials and extended affine Weyl groups of series A. *Ann. Inst. Fourier (Grenoble)* 53 (2003), no. 3, 927–940.
- [161] *T. Ohtsuki*, Quantum invariants. A study of knots, 3-manifolds, and their sets. *Series on Knots and Everything*, 29. World Scientific Publishing Co., Inc., River Edge, NJ, 2002. xiv+489 pp.
- [162] *E. Ossa*, On the cohomology of configuration spaces. *Algebraic topology: new trends in localization and periodicity (Sant Feliu de Guix, 1994)*, 353–361, *Progr. Math.*, 136, Birkh er, Basel, 1996.
- [163] *E. Ossa; V. V. Vershinin*, Thom spectra of commutants of generalized braid groups. *Funct. Anal. Appl.* 32 (1998), no. 4, 219–226 (1999), translated from Russian: *Funktsional. Anal. i Prilozhen.* 32 (1998), no. 4, 1–9.
- [164] *S. Yu. Orevkov*, Strong positivity in the right-invariant order on a braid group and quasipositivity. (Russian) *Mat. Zametki* 68 (2000), no. 5, 692–698; translation in *Math. Notes* 68 (2000), no. 5-6, 588–593.
- [165] *S. Yu. Orevkov*, Markov moves for quasipositive braids. *C. R. Acad. Sci. Paris Sr. I Math.* 331 (2000), no. 7, 557–562.
- [166] *S. Yu. Orevkov*, Quasipositivity test via unitary representations of braid groups and its applications to real algebraic curves. *J. Knot Theory Ramifications* 10 (2001), no. 7, 1005–1023.
- [167] *S. Yu. Orevkov*, Quasipositivity problem for 3-braids. *Turkish J. Math.* 28 (2004), no. 1, 89–93.

- [168] *S. Yu. Orevkov*, Solution of the word problem in the singular braid group. *Turkish J. Math.* 28 (2004), no. 1, 95–100.
- [169] *S. Yu. Orevkov; V. V. Shevchishin*, Markov theorem for transversal links. *J. Knot Theory Ramifications* 12 (2003), no. 7, 905–913.
- [170] *L. Paris*, Parabolic subgroups of Artin groups. *J. Algebra* 196 (1997), no. 2, 369–399.
- [171] *L. Paris*, Centralizers of parabolic subgroups of Artin groups of type A_l, B_l , and D_l . *J. Algebra* 196 (1997), no. 2, 400–435.
- [172] *L. Paris*, Artin monoids inject in their groups. *Comment. Math. Helv.* 77 (2002), no. 3, 609–637.
- [173] *L. Paris*, The proof of Birman’s conjecture on singular braid monoids. *Geom. Topol.* 8 (2004), 1281–1300 (electronic).
- [174] *L. Paris; D. Rolfsen*, Geometric subgroups of surface braid groups. *Ann. Inst. Fourier (Grenoble)* 49 (1999), no. 2, 417–472.
- [175] *V. V. Prasolov and A. B. Sossinsky*, Knots, links, braids and 3-manifolds. An introduction to the new invariants in low-dimensional topology. *Translations of Mathematical Monographs*, 154. American Mathematical Society, Providence, RI, 1997. viii+239 pp.
- [176] *K. Reidemeister*, *Knotentheorie*. (Erg. d. Math. u. ihrer Grenzgebiete. 1, No.1) Berlin: Julius Springer. VI, 74 S., 114 Fig. (1932). English translation: *Knot theory*. BCS Associates, Moscow, Idaho, 1983. xv+143 pp.
- [177] *A. G. Savushkina*, On the commutator subgroup of the braid group. (Russian) *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* 1993, , no. 6, 11–14, 118 (1994); translation in *Moscow Univ. Math. Bull.* 48 (1993), no. 6, 9–11.
- [178] *A. G. Savushkina*, On a group of conjugating automorphisms of a free group. (Russian) *Mat. Zametki* 60 (1996), no. 1, 92–108, 159; translation in *Math. Notes* 60 (1996), no. 1-2, 68–80 (1997).
- [179] *A. G. Savushkina*, The center of a colored braid group. (Russian) *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* 1996, , no. 1, 32–36, 103; translation in *Moscow Univ. Math. Bull.* 51 (1996), no. 1, 27–30.
- [180] *A. G. Savushkina*, Basis-conjugating automorphisms of a free group. (Russian) *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* 1996, , no. 4, 17–21, 110; translation in *Moscow Univ. Math. Bull.* 51 (1996), no. 4, 14–17.
- [181] *G. P. Scott*, Braid groups and the group of homeomorphisms of a surface. *Proc. Cambridge Philos. Soc.* 68, 1970, 605–617.
- [182] *V. Sergiescu*, Graphes planaires et présentations des groupes de tresses, *Math. Z.* 1993, 214, 477–490.
- [183] *A. B. Sossinsky*, Preparation theorems for isotopy invariants of links in 3-manifolds. *Quantum groups* (Leningrad, 1990), 354–362, *Lecture Notes in Math.*, 1510, Springer, Berlin, 1992.
- [184] *R. Street*, Braids among the groups. *Seminarberichte aus dem Fachbereich Mathematik*, Band 63, Teil 5 (1998) 699–703.
- [185] *I. Sysoeva*, Dimension n representations of the braid group on n strings. *J. Algebra* 243 (2001), no. 2, 518–538.
- [186] *P. Traczyk*, A new proof of Markov’s braid theorem. *Knot theory* (Warsaw, 1995), 409–419, *Banach Center Publ.*, 42, Polish Acad. Sci., Warsaw, 1998.
- [187] *V. Turaev*, Faithful linear representations of the braid groups. *Séminaire Bourbaki*, Vol. 1999/2000. *Astérisque* No. 276 (2002), 389–409.
- [188] *V. A. Vassiliev*, Complements of discriminants of smooth maps: topology and applications. *Translations of Mathematical Monographs*, 98. American Mathematical Society, Providence, RI, 1992. vi+208 pp.
- [189] *A. M. Vershik; S. Nechaev; R. Bikbov*, Statistical properties of locally free groups with applications to braid groups and growth of random heaps. *Comm. Math. Phys.* 212 (2000), no. 2, 469–501.
- [190] *V. V. Vershinin*, On braid groups in handlebodies. *Sib. Math. J.* 39, No.4, 645–654 (1998); translation from *Sib. Mat. Zh.* 39, No.4, 755–764 (1998).
- [191] *V. V. Vershinin*, On homological properties of singular braids. *Trans. Amer. Math. Soc.* 350 (1998), no. 6, 2431–2455.
- [192] *V. V. Vershinin*, Homology of braid groups and their generalizations. *Knot theory* (Warsaw, 1995), 421–446, *Banach Center Publ.*, 42, Polish Acad. Sci., Warsaw, 1998.

- [193] *V. V. Vershinin*, Generalizations of braids from a homological point of view. *Sib. Adv. Math.* 9, No.2, 109-139 (1999).
- [194] *V. V. Vershinin*, Mapping class groups and braid groups. *St. Petersburg Math. J.* 10 (1999), no. 6, 997-1003; translation from Russian: *Algebra i Analiz* 10 (1998), no. 6, 135-143.
- [195] *V. V. Vershinin*, Braid groups and loop spaces. *Russ. Math. Surv.* 54, No.2, 273-350 (1999); translation from *Usp. Mat. Nauk* 54, No.2, 3-84 (1999).
- [196] *V. V. Vershinin*, Thom spectra of generalized braid groups. *J. London Math. Soc.* (2) 61 (2000), no. 1, 245-258.
- [197] *V. V. Vershinin*, On homology of virtual braids and Burau representation. *Knots in Hellas '98*, Vol. 3 (Delphi). *J. Knot Theory Ramifications* 10 (2001), no. 5, 795-812.
- [198] *V. V. Vershinin*, On presentations of generalizations of braids with few generators, to appear in *Fundam. Prikl. Mat.*
- [199] *V. V. Vershinin*, On the singular braid monoid, [ArXiv:mathGR/0309339](https://arxiv.org/abs/math/0309339).
- [200] *J. Wu*, A braided simplicial group. *Proc. London Math. Soc.* (3) 84 (2002), no. 3, 645-662.
- [201] *O. Zariski*, On the Poincare group of rational plane curves, *Am. J. Math.* 1936, 58, 607-619.
- [202] *O. Zariski*, The topological discriminant group of a Riemann surface of genus p , *Am. J. Math.* 1937, 59, 335-358.
- [203] *J. Zhu*, On singular braids. *J. Knot Theory Ramifications*, 6 (1997), no. 3, 427-440.
- [204] *M. G. Zinno*, On Krammer's representation of the braid group. *Math. Ann.* 321 (2001), no. 1, 197-211.

DÉPARTEMENT DES SCIENCES MATHÉMATIQUES, UNIVERSITÉ MONTPELLIER II, PLACE EUGÈNE BATAILLON, 34095 MONTPELLIER CEDEX 5, FRANCE

E-mail address: `vershini@math.univ-montp2.fr`

SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK, 630090, RUSSIA

E-mail address: `versh@math.nsc.ru`